

FORM PTO-1390 (REV. 12-2001)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER <b>0380-P02825US0</b>	
<b>TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371</b>				U.S. APPLICATION NO. (If known, see 37 CFR 1.5 not yet assigned <b>10/088187</b>	
INTERNATIONAL APPLICATION NO. <b>PCT/GB00/03525</b>		INTERNATIONAL FILING DATE <b>13 September 2000</b>		PRIORITY DATE CLAIMED <b>17 September 1999</b>	
TITLE OF INVENTION <b>METHODS AND MEANS FOR MODIFICATION OF PLANT FLOWERING CHARACTERISTICS</b>					
APPLICANT(S) FOR DO/EO/US <b>DEAN, Caroline; LEVY, Yaron Yakov</b>					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below. 4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). b. <input checked="" type="checkbox"/> has been communicated by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). a. <input type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)). 9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).					
<b>Items 11 to 20 below concern document(s) or information included:</b>					
11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A <b>FIRST</b> preliminary amendment. 14. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment. 15. <input type="checkbox"/> A substitute specification. 16. <input type="checkbox"/> A change of power of attorney and/or address letter. 17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 20. <input checked="" type="checkbox"/> Other items or information: Copy of form PCT/IB/308					

U.S. APPLICATION NO. (if known, see 37 CFR 1.53) not yet assigned <b>70/0288187</b>		INTERNATIONAL APPLICATION NO. <b>PCT/GB00/03525</b>		ATTORNEY'S DOCKET NUMBER <b>0380-P02825US0</b>	
--	--	--	--	---	--

21. <input checked="" type="checkbox"/> The following fees are submitted: <b>BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO . . . . . <b>\$1040.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO . . . . . <b>\$890.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO . . . . . <b>\$740.00</b>  International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) . . . . . <b>\$710.00</b>  International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) . . . . . <b>\$100.00</b>  <b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b>				<b>CALCULATIONS PTO USE ONLY</b>          <table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:50%; text-align: right;">\$ 890.00</td> <td style="width:50%;"></td> </tr> </table>		\$ 890.00	
\$ 890.00							

Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
---	--	--	--	----	--

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$	
Total claims	34 - 20 =	14	x <b>\$18.00</b>	\$ 252.00	
Independent claims	4 - 3 =	1	x <b>\$84.00</b>	\$ 84.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ <b>\$280.00</b>	\$ 0	
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$ 1,226.00	

<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	
<b>SUBTOTAL =</b>				\$ 1,226.00	

Processing fee of <b>\$130.00</b> for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
<b>TOTAL NATIONAL FEE =</b>				\$ 1,226.00	

Fee for recording the enclosed assignment (37 CFR 1.21(h)) The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). <b>\$40.00</b> per property +				\$	
<b>TOTAL FEES ENCLOSED =</b>				\$ 1,226.00	

		Amount to be refunded:	\$
		charged:	\$

a. ☒ A check in the amount of \$ 1,226.00 to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_ to cover the above fees.  
 A duplicate copy of this sheet is enclosed.


c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any  
 overpayment to Deposit Account No. 04-1406. A duplicate copy of this sheet is enclosed.

d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card  
 information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR  
 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:  Kathleen D. Rigaut, Ph.D., J.D. DANN, DORFMAN, HERRELL AND SKILLMAN 1601 Market Street, Suite 720 Philadelphia, Pennsylvania 19103 United States of America	 SIGNATURE Kathleen D. Rigaut, Ph.D., J.D. NAME <u>43,047</u> REGISTRATION NUMBER
--	--

10/088187  
JC05 Rec'd PCT/PTO 15 MAR 2002

DANN, DORFMAN, HERRELL AND SKILLMAN

A PROFESSIONAL CORPORATION

1601 MARKET STREET · SUITE 720 · PHILADELPHIA, PA · 19103-2307

PHONE (215) 563-4100 · FAX (215) 563-4044

COMMISSIONER OF PATENTS AND TRADEMARKS WASHINGTON, DC 20231	DATE OF MAILING  15 March 2002
BOX PCT	APPLICANT'S OR AGENT'S FILE REF. 0380-P02825US0
IDENTIFICATION OF THE INTERNATIONAL APPLICATION	
INTERNATIONAL APPLICATION NO. PCT/GB00/03525	INTERNATIONAL FILING DATE 13 September 2000
APPLICANT (name) PLANT BIOSCIENCE LIMITED	
TRANSMITTAL OF LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US)	

CERTIFICATE OF MAILING BY EXPRESS MAIL UNDER 37 C.F.R. §1.10

NO. OF EXPRESS MAIL LABEL EL680721249US

DATE OF DEPOSIT WITH POSTAL SERVICE March 15, 2002

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. §1.10 on the date indicated above, and is addressed to the COMMISSIONER OF PATENTS, BOX PCT, Washington, DC 20231-0001.

Jane C. Bogan  
Name of Person Mailing Paper or Fee

Jane C. Bogan  
Signature of Person Mailing Paper or Fee

THE UNITED STATES PATENT AND TRADEMARK OFFICE

United States Serial No. : Not yet assigned  
International Application No. : PCT/GB00/03525  
International Filing Date : 13 September 2000  
Inventor(s) : Caroline Dean et al.  
Title : METHODS AND MEANS FOR  
MODIFICATION OF PLANT  
FLOWERING CHARACTERISTICS

-----  
Suite 720  
1601 Market Street  
Philadelphia, PA 19103-2307  
(215) 563-4100 (telephone)  
(215) 563-4044 (facsimile)  
Our File: 0380-P02825US0

---

Box PCT  
Assistant Commissioner  
for Patents  
Washington, D.C. 20231

**PRELIMINARY AMENDMENT**

Dear Sir:

Before calculation of the filing fee, please amend the  
above-identified application as follows:

In the specification:

At page 1, line 3, please insert the following priority  
claim:

-- This application is a §371 filing of PCT/GB00/03525 filed  
13 September 2000, which in turn claims priority to GB  
application 9922071.7 filed 17 September 1999, the entire  
disclosures of each being incorporated by reference herein.--

At page 80 of the specification, please insert a copy of the  
abstract of the specification which is attached hereto on a  
separate sheet.

In the claims:

Please amend claims 3-6, 8, 11-14, 16, 18-20, 22, 24-25, 27, 29-30 and 32-33 of the above-referenced patent application, which claims are based on the Article 34 claim amendments filed in the corresponding international patent application, as follows:

3. (Amended) A nucleic acid as claimed in claim 1 wherein the VRN1 nucleotide sequence is that shown in Fig 7 from nucleotides 269-1295 inclusive, or a sequence which is degeneratively equivalent thereto.
4. (Amended) A nucleic acid as claimed in claim 1 wherein the VRN1 nucleotide sequence is shown in Annex I.
5. (Amended) A nucleic acid as claimed in claim 1 wherein the VRN1 nucleotide sequence encodes a derivative of the polypeptide shown in Fig 7 by way of addition, insertion, deletion or substitution of one or more amino acids.
6. (Amended) A nucleic acid as claimed in claim 1 wherein the VRN1 nucleotide sequence consists of an allelic or other homologous variant of the nucleotide sequence wherein the VRN1 nucleotide sequence is that shown in Fig 7 from nucleotides 269-1295 inclusive, or a sequence which is degeneratively equivalent thereto.
8. (Amended) An isolated nucleic acid which comprises a nucleotide sequence which is the complement of the VRN1 nucleotide sequence of claim 1.
11. (Amended) A process for producing a nucleic acid comprising the VRN1 nucleotide sequence encoding a derivative of the polypeptide shown in Fig 7 by way of addition, insertion, deletion or substitution of one or more amino acids or sequence degeneratively equivalent thereto.

13. (Amended) A method as claimed in claim 12, which method comprises the steps of:
  - a. providing a preparation of nucleic acid from a plant cell;
  - b. providing said probe or primer sequence;
  - c. contacting nucleic acid in said preparation with said probe or primer sequence under conditions for hybridisation; and,
  - d. identifying nucleic acid in said preparation which hybridises with said nucleic acid molecule.
14. (Amended) A method as claimed in claim 12, which method comprises the steps of:
  - a. providing a preparation of nucleic acid from a plant cell;
  - b. providing a pair of said primers, said primers being suitable for PCR;
  - c. contacting nucleic acid in said preparation with said primers under conditions for performance of PCR;
  - d. performing PCR and determining the presence or absence of an amplified PCR product.



vernalisation response of plant into which the nucleic acid is introduced and expressed, in a suitable host cell.

29. (Amended) A method for assessing the vernalisation phenotype of a plant, the method comprising the step of determining the presence and/or identity of a VRN1 allele therein comprising the use of a nucleic acid selected from the group consisting of a probe or primer having a sequence of about 16-24 nucleotides in length present in Annex I, a complementary sequence of a sequence present in Annex I, a sequence degeneratively equivalent to a sequence present in Annex I, and a sequence as claimed in claim 10.
30. (Amended) A method for influencing or affecting the vernalisation phenotype of a plant, which method comprises the step of causing or allowing expression of a heterologous nucleic acid as claimed in claim 1 within the cells of the plant, following an earlier step of introducing the nucleic acid into a cell of the plant or an ancestor thereof.
32. (Amended) A method as claimed in claim 30 for reducing the vernalisation requirement of a plant, wherein the heterologous nucleic acid is a VRN1 nucleotide sequence encoding a polypeptide which is capable of specifically altering the vernalisation response of plant into which the nucleic acid is introduced and expressed.
33. (Amended) A method as claimed in claim 30 for increasing the vernalisation requirement of a plant, which method comprises any of the following steps of:
  - (i) causing or allowing transcription from a nucleic acid which is the complement of a VRN1 sequence in the plant such as to reduce VRN1 expression by an antisense mechanism;



- \*\*\*\*\*
- (ii) causing or allowing transcription from a nucleic acid which is a VRN1 nucleotide sequence encoding a polypeptide which is capable of specifically altering the vernalisation response of plant into which the nucleic acid is introduced and expressed such as to reduce VRN1 expression by co-suppression;
  - (iii) use of nucleic acid encoding a ribozyme specific for a nucleic acid selected from the group consisting of a VRN1 sequence which encodes the VRN1 polypeptide of Fig. 7 and a variant resistance polypeptide shown in Fig. 7 which shares at least about 50%, 60%, 70%, 80% or 90% identity therewith.


**REMARKS**

The purpose of this Preliminary Amendment is to 1) eliminate multiple claims dependencies and to eliminate certain claims which, due to their form, do not comply with current United States Patent and Trademark Office practice; 2) provide an abstract of the disclosure on a separate sheet; and 3) insert a proper priority claim into the application.

Entry of the foregoing amendments is respectfully requested, as they do not introduce new matter into the present application, and, therefore should not be deemed objectionable.

Early and favorable consideration of the present application is respectfully requested.

Respectfully submitted,

  
Kathleen D. Rigant, Ph.D., J.D.  
Reg. No. 43,047  
Attorney for Applicant

PJH:KDR:ksk

**MARKED-UP COPY OF THE CLAIMS**

3. (Amended) A nucleic acid as claimed in claim 1 [or claim 2] wherein the VRN1 nucleotide sequence is that shown in Fig 7 from nucleotides 269-1295 inclusive, or a sequence which is degeneratively equivalent thereto.
4. (Amended) A nucleic acid as claimed in claim 1 [or claim 2] wherein the VRN1 nucleotide sequence is shown in Annex I.
5. (Amended) A nucleic acid as claimed in claim 1 [or claim 2] wherein the VRN1 nucleotide sequence encodes a derivative of the polypeptide shown in Fig 7 by way of addition, insertion, deletion or substitution of one or more amino acids.
6. (Amended) A nucleic acid as claimed in claim 1 [or claim 2] wherein the VRN1 nucleotide sequence consists of an allelic or other homologous variant of the nucleotide sequence [of claim 3] wherein the VRN1 nucleotide sequence is that shown in Fig 7 from nucleotides 269-1295 inclusive, or a sequence which is degeneratively equivalent thereto.
8. (Amended) An isolated nucleic acid which comprises a nucleotide sequence which is the complement of the VRN1 nucleotide sequence of [any one of the preceding] claim[s] 1.
11. (Amended) A process for producing a nucleic acid [as claimed in claim 5] comprising the VRN1 nucleotide sequence encoding a derivative of the polypeptide shown in Fig 7 by way of addition, insertion, deletion or substitution of one or more amino acids or sequence degeneratively equivalent thereto [comprising the step of modifying a nucleic acid as claimed in claim 3 or claim 4].



claimed in claim 9 or claim 10];

- c. contacting nucleic acid in said preparation with said primers under conditions for performance of PCR;
  - d. performing PCR and determining the presence or absence of an amplified PCR product.
16. (Amended) A recombinant vector which comprises the nucleic acid of [any one of] claim[s] 1 [to 8].
  18. (Amended) A vector as claimed in [claim 16 or] claim 17 which is a plant vector.
  19. (Amended) A method for transforming a host cell, which comprises the step of introducing the vector of [any one of claim[s] 16 to] 18 into a host cell, and optionally causing or allowing recombination between the vector and the host cell genome such as to transform the host cell.
  20. (Amended) A host cell containing or transformed with a heterologous vector of [any one of] claim[s] 16 to] 18.
  22. (Amended) A transgenic plant which is obtainable by the method of claim 21, or which is a clone, or selfed or hybrid progeny or other descendant of said transgenic plant, which in each case includes a heterologous nucleic acid [of any one of claims 1 to 8] wherein said heterologous nucleic acid is a VRN1 nucleotide sequence encoding a polypeptide which is capable of specifically altering the vernalisation response of plant into which the nucleic acid is introduced and expressed or a variant thereof.
  24. (Amended) A part of propagule from a plant as claimed in claim 22 [or claim 23, which in either case includes a heterologous nucleic acid of any one of claims 1 to 8].

25. (Amended) An isolated polypeptide which is encoded by the VRN1 nucleotide sequence of [any one of] claim[s] 1 [to 7].
27. (Amended) A method of making the polypeptide of claim [25 or claim] 26, which method comprises the step of causing or allowing expression from a [nucleic acid] VRN1 nucleotide sequence encoding a polypeptide which is capable of specifically altering the vernalisation response of plant into which the nucleic acid is introduced and expressed, [of any one of claims 1 to 7] in a suitable host cell.
29. (Amended) A method for assessing the vernalisation phenotype of a plant, the method comprising the step of determining the presence and/or identity of a VRN1 allele therein comprising the use of a nucleic acid [as claimed in claim 9 or] selected from the group consisting of a probe or primer having a sequence of about 16-24 nucleotides in length present in Annex I, a complementary sequence of a sequence present in Annex I, a sequence degeneratively equivalent to a sequence present in Annex I, and a sequence as claimed in claim 10.
30. (Amended) A method for influencing or affecting the vernalisation phenotype of a plant, which method comprises the step of causing or allowing expression of a heterologous nucleic acid as claimed in [any one of] claim[s] 1 [to 8] within the cells of the plant, following an earlier step of introducing the nucleic acid into a cell of the plant or an ancestor thereof.
32. (Amended) A method as claimed in claim 30 [or claim 31] for reducing the vernalisation requirement of a plant, wherein the heterologous nucleic acid is [that claimed in any one of claims 1 to 7] a VRN1 nucleotide sequence encoding a polypeptide which is capable of specifically altering the

vernalisation response of plant into which the nucleic acid is introduced and expressed.

33. (Amended) A method as claimed in claim 30 [or claim 31] for increasing the vernalisation requirement of a plant, which method comprises any of the following steps of:
- (i) causing or allowing transcription from a nucleic acid [as claimed in claim 8]which is the complement of a VRN1 sequence in the plant such as to reduce VRN1 expression by an antisense mechanism;
  - (ii) causing or allowing transcription from a nucleic acid [as claimed in any one of claims 1 to 7 or a part thereof] which is a VRN1 nucleotide sequence encoding a polypeptide which is capable of specifically altering the vernalisation response of plant into which the nucleic acid is introduced and expressed such as to reduce VRN1 expression by co-suppression;
  - (iii) use of nucleic acid encoding a ribozyme specific for a nucleic acid [as claimed in any one of claims 1 to 7] selected from the group consisting of a VRN1 sequence which encodes the VRN1 polypeptide of Fig. 7 and a variant resistance polypeptide shown in Fig. 7 which shares at least about 50%, 60%, 70%, 80% or 90% identity therewith.

## ABSTRACT

Provided are isolated nucleic acid molecules which comprise VRN1 nucleotide sequences, which encode a polypeptide which is capable of specifically altering the vernalisation response of a plant into which the nucleic acid is introduced and expressed. Examples include cDNA and gDNA sequences (see e.g., Annex I). Also provided are variant molecules which may be derivatives or homologues (e.g., alleles, or paralogues such as *RTV1*), plus also complementary molecules. Corresponding polypeptides form a further part of the invention. The invention also provides methods and materials for preparing and using these molecules, e.g., in the production of plants having modified vernalisation characteristics. Also provided are methods for influencing and assessing the vernalisation phenotype of a plant.

METHODS AND MEANS FOR MODIFICATION OF PLANT FLOWERINGCHARACTERISTICSTECHNICAL FIELD

5

The present invention relates generally to methods and materials for use in modifying plant characteristics, particularly the vernalization response in plants.

10 PRIOR ART

Plants must integrate a wide variety of environmental signals in order to maximize their reproductive success. Part of this integration must involve perception of the seasons, both to  
15 ensure the plant flowers during the correct season (for which it is adapted) and to synchronise its flowering with other members of its own species, to increase the chances of cross-fertilization. *Arabidopsis thaliana* serves as a model plant, for it exhibits responses to a wide variety of environmental  
20 stimuli that are observed in many species. Amongst other stimuli, flowering in naturally occurring strains (ecotypes) of *Arabidopsis* can be promoted by vernalization, a long cold treatment that mimics the cold of winter.

25 Many species of plants that grow in temperate or cooler climes have an obligate requirement for vernalization in order to flower. Such plants typically germinate in autumn, and over winter as vegetative plants, and flower in milder conditions of spring. Vernalization thus acts as a cue, to allow plants  
30 to sense the seasons, and to time their flowering to maximise their chance of reproductive success.

Species for which flowering is important to crop production are numerous, essentially all crops which are grown from seed,  
35 with important examples being the cereals, rice and maize,



probably the most agronomically important in warmer climatic zones, and wheat, barley, oats and rye in more temperate climates. Important seed products are oil seed rape, sugar beet, maize, sunflower, soybean and sorghum. Many crops which are harvested for their roots are, of course, grown annually from seed and the production of seed of any kind is very dependent upon the ability of the plant to flower, to be pollinated and to set seed. In horticulture, control of the timing of flowering is important. Horticultural plants whose flowering may be controlled include lettuce, endive and vegetable brassicas including cabbage, broccoli and cauliflower, and carnations and geraniums.

In view of the large number of commercially important crop species which have a requirement for vernalization in order to flower, modification of this requirement (e.g. by reducing the duration of vernalization required, or changing the optimum temperature, or abrogating the requirement altogether) would be of agronomic interest.

#### DISCLOSURE OF THE INVENTION

The inventors have used a late flowering, vernalization responsive mutant of *Arabidopsis*, the *fca* mutant, as a background in which to isolate mutants that exhibit a reduced vernalization response and to identify *vrn1* alleles which are responsible for this phenotype. The *VRN1* gene is the first *Arabidopsis* flowering time gene to be isolated that is apparently exclusive to the vernalization promotion pathway. As discussed in more detail below, manipulation of the gene may permit the control or modification of the vernalization response of agronomically important crop species.

That *VRN1* is required for a normal vernalization response is clear from the phenotype of the *vrn1* mutants. Further

experiments by the inventors indicate that there is a quantitative aspect to VRN1 activity. This suggests that artificially increasing or decreasing the amount of VRN1 (e.g., through overexpression or antisense suppression) may provide a tool to, *inter alia* fine-tune the kinetics and/or optimal temperature of the vernalization response; render plants immune to the effect of cold on flowering response; or alleviate the requirement for cold treatment altogether. In addition to quantitative manipulation, a further layer of control could be obtained by driving VRN1 sense or anti-sense constructs using promoters that are either on all the time (constitutive); inducible upon application of a specific molecule; or which "naturally" drive expression only during a certain portion of the plant life cycle, e.g., seed maturation or late vegetative phase.

Such methods could be used to improve agronomically important crop species, for instance as follows:

- (a) Extension of geographic range of elite varieties: If an elite cultivar of a crop originates from a geographic area where it has adapted to require a certain vernalization period, and it is therefore climatically-limited in its range, then fine-tuning the expression of VRN1 may permit alteration of the length and intensity of cold treatment required to achieve an optimum flowering time in new geographic areas. Two facts are noteworthy in this regard: (1) even modest alterations in vernalization response could open up huge new areas of cultivation for particular elite varieties (an analogous situation to that in which small changes in climatic conditions can alter the ecology and character of huge areas of landscape), and (2) the commercial success of elite genotypes is largely hampered by dependency on specific climatic conditions found in limited geographic areas.

(b) Shortening of vernalization period: if a winter crop can be sown and left in the ground for a shorter period than usual (i.e. a reduced vernalization time, perhaps resulting from increased- or mis-expression of *VRN1*) this may reduce the risk associated with severe winter weather conditions, as the crops are exposed to winter conditions for a shorter time.

(c) Extension of vegetative growth: If the crop in question is one in which the vegetative portions of the plant are the desired product (e.g., leaf vegetables, sugar beet), then preventing the plant from flowering in response to cold temperature (i.e., by rendering it less sensitive to the cold by impairing *VRN1* function) would prevent diversion of valuable plant resources from the vegetative tissues to the developing reproductive tissues, thereby increasing yield.

Further experiments indicate that species other than *Arabidopsis* contain genes similar to *VRN1*. Additionally, homologues and/or orthologues and/or paralogues of *VRN1* (such as *RTV1*) may also exist in *Arabidopsis* and other species. Based on the disclosure herein, such genes may be isolated without undue burden by those skilled in the art and used analogously to those disclosed herein.

These and other aspects of the present invention will now be discussed in more detail.

Thus according to one aspect of the present invention there is provided an isolated nucleic acid molecule which is capable of specifically altering the vernalisation response of a plant into which the nucleic acid is introduced.

The alteration in the vernalisation response may be assessed by comparison with a plant in which the nucleic acid has not been so introduced.

The vernalization response phenotype of plants may be investigated by examining their flowering time in response to differing durations of vernalization treatment. In the experiments below this was assessed in two ways: (1) as the total number of vegetative leaves produced prior to flowering (LN), and (2) as the time in days from the end of the vernalization treatment to the appearance of the first floral bud (FT). However any appropriate method known to those skilled in the art may be used.

Apart from the specific change in vernalisation response, it is preferred that other characteristics of the plant are substantially unchanged by the polypeptide, which is to say that the polypeptide acts specifically on this response and not more generally on flowering time characteristics or other stimuli, such as those mediated by other loci such as the *FRI* locus (Clarke and Dean, 1994, Mol. Gen. Genet. 242, 81-89) or the *VRN2* locus (Chandler et al., 1996).

Preferably the isolated nucleic acid molecule capable of specifically altering the vernalisation response of a plant is obtainable from the *VRN1* locus of a plant, more preferably from *A. thaliana*.

Nucleic acid according to the present invention may include cDNA, RNA, genomic DNA and modified nucleic acids or nucleic acid analogs (e.g. peptide nucleic acid). Where a DNA sequence is specified, e.g. with reference to a figure, unless context requires otherwise the RNA equivalent, with U substituted for T where it occurs, is encompassed. Nucleic acid molecules according to the present invention may be provided isolated and/or purified from their natural environment, in substantially pure or homogeneous form, or free or substantially free of other nucleic acids of the species of origin. Where used herein, the term "isolated" encompasses

all of these possibilities. The nucleic acid molecules may be wholly or partially synthetic. In particular they may be recombinant in that nucleic acid sequences which are not found together in nature (do not run contiguously) have been ligated  
5 or otherwise combined artificially. Alternatively they may have been synthesised directly e.g. using an automated synthesiser.

Thus in one aspect of the invention, there is disclosed a  
10 nucleic acid encoding the polypeptide of Fig 7. The VRN1 polypeptide is 341 amino acids in length and is comprised of at least three regions. Region 1 (residues 2-94 in Figure 7) and 3 (residues 239-332) can be aligned to each other, and are related to the B3 DNA-binding domain originally found in the  
15 maize transcription factor VIVIPAROUS1 (VP1; McCarty et al., 1991). Region 2 of VRN1 (residues 95-238), which lies between the two B3 domains is not obviously related to any domain of known function, nor does it have obvious features of a transcriptional activation or repression domain. Nonetheless,  
20 region 2 does contain several provocative sequence features and motifs, including a putative nuclear localization signal (NSL), two putative PEST regions, and three RXXL motifs also associated with rapid protein degradation (Cooper et al., 1997). Interestingly, the second PEST region of VRN1 contains  
25 a potential protein kinase C (PKC) phosphorylation site (residues 176-178).

One nucleic acid encoding this polypeptide is shown in Fig 7 from nucleotides 269-1295 inclusive (including stop codon).

30 Other nucleic acids of the invention include those which are degeneratively equivalent to this.

A genomic sequence including the VRN1 locus is shown in Annex I. The putative cDNA sequence transcribed from this genomic  
35 sequence is shown at Fig 7. Although this ORF has been

designated the VRN1 ORF herein, it will be appreciated by those skilled in the art that the discussion hereinafter applies equally to any other ORF present in the described sequence which has the properties attributed to VRN1.

5

In a further aspect of the present invention there are disclosed nucleic acids which are variants of the VRN1 sequences discussed above.

- 10 A variant nucleic acid molecule shares homology with, or is identical to, all or part of the sequences discussed above.

Such variants may be used to alter the vernalisation characteristics of a plant, as assessed by the methods

- 15 disclosed herein. For instance a variant nucleic acids may be include a sequence encoding a functional polypeptide (e.g. which may be a variant of the VRN1 polypeptide and which may cross-react with an antibody raised to said polypeptide). Alternatively they may include a sequence which interferes  
20 with the expression or activity of such a polypeptide (e.g. sense or anti-sense suppression of a VRN1 coding sequence).

- Variants may also be used to isolate or amplify nucleic acids which have these properties (e.g. by inclusion of a sequence  
25 which is hybridisable with a VRN1 sequence.).

Generally speaking variants may be:

- (i) Novel, naturally occurring, nucleic acids, isolatable  
30 using the sequences of the present invention. They may include alleles (which will include polymorphisms or mutations at one or more bases - for instance *vrn1-1* or *vrn1-2* shown in Fig 7) or pseudoalleles (which may occur at closely linked loci to the VRN1 gene). Also included are paralogues,  
35 isogenes, or other homologous genes belonging to the same

family as the VRN1 gene. Although these may occur at different genomic loci to the gene, they are likely to share conserved regions with it (see e.g. RTV1 in the Examples below). Also included are homologues of VRN1 from other plant species.

(ii) Artificial nucleic acids, which can be prepared by the skilled person in the light of the present disclosure. Such derivatives may be prepared, for instance, by site directed or random mutagenesis, or by direct synthesis. Preferably the variant nucleic acid is generated either directly or indirectly (e.g. via one or more amplification or replication steps) from an original nucleic acid having all or part of the VRN1 sequence shown in Fig 7.

Particularly included are variants which comprise only a distinctive part or fragment (however produced) corresponding to a portion of the sequence provided. The fragments may encode particular functional parts of the polypeptide.

Alternatively, the fragments may have utility in probing for, or amplifying, the sequence provided or closely related ones. Suitable lengths of fragment, and conditions, for such processes are discussed in more detail below.

Also included are nucleic acids corresponding to those above, but which have been extended at the 3' or 5' terminus.

The term 'variant' nucleic acid as used herein encompasses all of these possibilities. When used in the context of polypeptides or proteins it indicates the encoded expression product of the variant nucleic acid.

Some of the aspects of the present invention relating to variants will now be discussed in more detail.

Homology (similarity or identity) may be assessed as set out in the Materials and Methods section in the Examples below.

Homology may be at the nucleotide sequence and/or encoded amino acid sequence level. Preferably, the nucleic acid and/or amino acid sequence shares at least about 65%, or 70%, or 80% identity, most preferably at least about 90%, 95%, 96%, 97%, 98% or 99% identity.

Homology may be over the full-length of the relevant sequence shown herein, or may be over a part of it, preferably over a contiguous sequence of about or greater than about 20, 25, 30, 33, 40, 50, 67, 133, 167, 200, 233, 267, 300, or more amino acids or codons, compared with Fig 7.

15

Thus a variant polypeptide encoded by a nucleic acid of the present invention may include within the sequence shown in Fig 7, a single amino acid or 2, 3, 4, 5, 6, 7, 8, or 9 changes, about 10, 15, 20, 30, 40 or 50 changes, or greater than about 50, 60, 70, 80 or 90 changes.

20

In a further aspect of the invention there is disclosed a method of producing a derivative nucleic acid comprising the step of modifying any of the sequences disclosed above, particularly the coding sequence of Fig 7.

25

Changes may be desirable for a number of reasons. For instance they may introduce or remove restriction endonuclease sites or alter codon usage.

30

Alternatively changes to a sequence may produce a derivative by way of one or more of addition, insertion, deletion or substitution of one or more nucleotides in the nucleic acid, leading to the addition, insertion, deletion or substitution of one or more amino acids in the encoded polypeptide.

35



Such changes may modify sites which are required for post translation modification such as cleavage sites in the encoded polypeptide; motifs in the encoded polypeptide for phosphorylation etc. (e.g. residues 176-178 in Figure 7).

- 5 Leader or other targeting sequences (e.g. membrane or golgi locating sequences) may be added to the expressed protein to determine its location following expression if it is desired to isolate it from a microbial system.
- 10 Other desirable mutations may be random or site directed mutagenesis in order to alter the activity (e.g. specificity) or stability of the encoded polypeptide. Changes may be by way of conservative variation, i.e. substitution of one hydrophobic residue such as isoleucine, valine, leucine or
- 15 methionine for another, or the substitution of one polar residue for another, such as arginine for lysine, glutamic for aspartic acid, or glutamine for asparagine. As is well known to those skilled in the art, altering the primary structure of a polypeptide by a conservative substitution may not
- 20 significantly alter the activity of that peptide because the side-chain of the amino acid which is inserted into the sequence may be able to form similar bonds and contacts as the side chain of the amino acid which has been substituted out. This is so even when the substitution is in a region which is
- 25 critical in determining the peptides conformation. Also included are variants having non-conservative substitutions. As is well known to those skilled in the art, substitutions to regions of a peptide which are not critical in determining its conformation may not greatly affect its activity because they
- 30 do not greatly alter the peptide's three dimensional structure. In regions which are critical in determining the peptides conformation or activity such changes may confer advantageous properties on the polypeptide. Indeed, changes such as those described above may confer slightly advantageous
- 35 properties on the peptide e.g. altered stability or

specificity.

Particular regions, or domains, of VRN1 may have utility in their own right. For instance the B3 domains may be used to  
5 direct gene expression in a precise manner, for instance by the recognition of specific DNA sequences that represent elements in the promoters of their normal target genes. By creating fusion proteins, comprising the DNA binding domain (or domains) of VRN1, and a heterologous activation or  
10 repression domain borrowed from another protein, the expression of VRN1 target genes could be controlled. This may lead to a precise control of the expression of those genes that are normally targets of VRN1. Given that such genes, alone or in combination, ultimately control the transition to  
15 flowering (usually following vernalization), their directed expression in other conditions may provide a useful means to control flowering. Furthermore, the use of fusions based on the DNA binding domains in conventional SELEX or one-hybrid experiments may be used to reveal the target genes or DNA  
20 sequences normally bound by VRN1. Thus nucleic acids encoding these domains, or fusion proteins comprising them, form one embodiment of this aspect of the present invention.

In a further aspect of the present invention there is provided  
25 a method of identifying and/or cloning a VRN1 nucleic acid variant from a plant which method employs a sequence described above.

In one embodiment, nucleotide sequence information provided  
30 herein may be used in a data-base (e.g. of ESTs, or STSs) search to find homologous sequences, such as those which may become available in due course, and expression products of which can be tested for activity as described below.

35 For example, searches were conducted using the tBLASTn Program

(version 2.0 using the default parameters) available from NCBI (website: [www.ncbi.nlm.nih.gov/blast/](http://www.ncbi.nlm.nih.gov/blast/)). The 341 amino acid deduced VRN1 protein sequence was searched against all GenBank ESTs (dbEST database). Accessions are listed below which

5 satisfied the following criteria: (1) they were expressed - all sequences are ESTs (partial), i.e., derived from mRNA; (2) they shared VRN1 domain structure - all sequences share homology with VRN1 that extends beyond either of the two B3 domains, i.e., they are not simply one of many B3-containing

10 sequences; (3) The partial sequences share greater than or equal to 50% identity with VRN1 at the encoded amino acid level. In the light of the present invention, these partial sequences may be expected to be derived from hitherto uncharacterised VRN1-related genes.

15

	<u>Species</u>	<u>GenBank accession:</u>	<u>%ID</u>	<u>%similarity</u>
	Medicago truncatula	AW686695	86	91
	Medicago truncatula	AW584452	82	91
20	Glycine max	AW705298	75	86
	Gossypium hirsutum	AW187216	74	82
	Medicago truncatula	AW586752	72	84
	Gossypium arboreum	AW668373	69	85
	Gossypium arboreum	BE052554	66	82
25	Gossypium arboreum	BE054829	66	78
	Medicago truncatula	AW685743	54	59
	Medicago truncatula	AW685178	53	62
	Medicago truncatula	BE203124	51	67
	Medicago truncatula	AW736517	51	65

30

(Medicago truncatula=barrel medic, Glycine max=soybean, Gossypium hirsutum=cotton, Gossypium arboreum=tree cotton).

In another embodiment the nucleotide sequence information

35 provided herein may be used to design probes and primers for

probing or amplification. An oligonucleotide for use in  
probing or PCR may be about 30 or fewer nucleotides in length  
(e.g. 18, 21 or 24). Generally specific primers are upwards  
of 14 nucleotides in length. For optimum specificity and cost  
5 effectiveness, primers of 16-24 nucleotides in length may be  
preferred. Those skilled in the art are well versed in the  
design of primers for use in processes such as PCR. If  
required, probing can be done with entire restriction  
fragments of the gene disclosed herein which may be 100's or  
10 even 1000's of nucleotides in length. Small variations may be  
introduced into the sequence to produce 'consensus' or  
'degenerate' primers if required.

Such probes and primers form one aspect of the present  
15 invention.

Probing may employ the standard Southern blotting technique.  
For instance DNA may be extracted from cells and digested with  
different restriction enzymes. Restriction fragments may then  
20 be separated by electrophoresis on an agarose gel, before  
denaturation and transfer to a nitrocellulose filter.  
Labelled probe may be hybridised to the single stranded DNA  
fragments on the filter and binding determined. DNA for  
probing may be prepared from RNA preparations from cells.  
25 Probing may optionally be done by means of so-called 'nucleic  
acid chips' (see Marshall & Hodgson (1998) Nature  
Biotechnology 16: 27-31, for a review).

In one embodiment, a variant in accordance with the present  
30 invention is obtainable by means of a method which includes:

(a) providing a preparation of nucleic acid, e.g. from plant  
cells. Test nucleic acid may be provided from a cell as  
genomic DNA, cDNA or RNA, or a mixture of any of these,  
35 preferably as a library in a suitable vector. If genomic DNA

is used the probe may be used to identify untranscribed regions of the gene (e.g. promoters etc.), such as are described hereinafter,

- (b) providing a nucleic acid molecule which is a probe or  
5 primer as discussed above,
- (c) contacting nucleic acid in said preparation with said nucleic acid molecule under conditions for hybridisation of said nucleic acid molecule to any said gene or homologue in said preparation, and,
- 10 (d) identifying said gene or homologue if present by its hybridisation with said nucleic acid molecule. Binding of a probe to target nucleic acid (e.g. DNA) may be measured using any of a variety of techniques at the disposal of those skilled in the art. For instance, probes may be  
15 radioactively, fluorescently or enzymatically labelled. Other methods not employing labelling of probe include amplification using PCR (see below), RN'ase cleavage and allele specific oligonucleotide probing. The identification of successful hybridisation is followed by isolation of the nucleic acid  
20 which has hybridised, which may involve one or more steps of PCR or amplification of a vector in a suitable host.

Preliminary experiments may be performed by hybridising under low stringency conditions. For probing, preferred conditions  
25 are those which are stringent enough for there to be a simple pattern with a small number of hybridisations identified as positive which can be investigated further.

For example, hybridizations may be performed, according to the  
30 method of Sambrook et al. (below) using a hybridization solution comprising: 5X SSC (wherein 'SSC' = 0.15 M sodium chloride; 0.15 M sodium citrate; pH 7), 5X Denhardt's reagent, 0.5-1.0% SDS, 100 µg/ml denatured, fragmented salmon sperm DNA, 0.05% sodium pyrophosphate and up to 50% formamide.  
35 Hybridization is carried out at 37-42°C for at least six hours.

Following hybridization, filters are washed as follows: (1) 5 minutes at room temperature in 2X SSC and 1% SDS; (2) 15 minutes at room temperature in 2X SSC and 0.1% SDS; (3) 30 minutes - 1 hour at 37°C in 1X SSC and 1% SDS; (4) 2 hours at 42-65°C in 1X SSC and 1% SDS, changing the solution every 30 minutes.

One common formula for calculating the stringency conditions required to achieve hybridization between nucleic acid molecules of a specified sequence homology is (Sambrook et al., 1989):

$$T_m = 81.5^{\circ}\text{C} + 16.6\text{Log} [\text{Na}^+] + 0.41 (\% \text{ G+C}) - 0.63 (\% \text{ formamide}) - 600/\text{\#bp in duplex}$$

As an illustration of the above formula, using  $[\text{Na}^+] = [0.368]$  and 50-% formamide, with GC content of 42% and an average probe size of 200 bases, the  $T_m$  is 57°C. The  $T_m$  of a DNA duplex decreases by 1 - 1.5°C with every 1% decrease in homology. Thus, targets with greater than about 75% sequence identity would be observed using a hybridization temperature of 42°C. Such a sequence would be considered substantially homologous to the nucleic acid sequence of the present invention.

It is well known in the art to increase stringency of hybridisation gradually until only a few positive clones remain. Other suitable conditions include, e.g. for detection of sequences that are about 80-90% identical, hybridization overnight at 42°C in 0.25M  $\text{Na}_2\text{HPO}_4$ , pH 7.2, 6.5% SDS, 10% dextran sulfate and a final wash at 55°C in 0.1X SSC, 0.1% SDS. For detection of sequences that are greater than about 90% identical, suitable conditions include hybridization overnight at 65°C in 0.25M  $\text{Na}_2\text{HPO}_4$ , pH 7.2, 6.5% SDS, 10% dextran sulfate and a final wash at 60°C in 0.1X SSC, 0.1% SDS.

Thus this aspect of the present invention includes a nucleic acid including or consisting essentially of a nucleotide sequence of complementary to a nucleotide sequence

5 hybridisable with any encoding sequence provided herein. Another way of looking at this would be for nucleic acid according to this aspect to be hybridisable with a nucleotide sequence complementary to any encoding sequence provided herein.

10

In a further embodiment, hybridisation of nucleic acid molecule to a variant may be determined or identified indirectly, e.g. using a nucleic acid amplification reaction, particularly the polymerase chain reaction (PCR). PCR

15 requires the use of two primers to specifically amplify target nucleic acid, so preferably two nucleic acid molecules with sequences characteristic of VRN1 are employed. Using RACE PCR, only one such primer may be needed (see "PCR protocols; A Guide to Methods and Applications", Eds. Innis et al, Academic  
20 Press, New York, (1990)).

Preferred primers for amplification of conserved regions of VRN1 for use as probes to obtain genomic or cDNA clones may include any of those shown in Table 3.

25

For instance primers S63 and S49 may be used to amplify a VRN1 genomic region including the promoter and 3' end of the gene.

Primers V7 and V2 amplify the VRN1 cDNA ORF. Primers V6 and  
30 V15 may be used to distinguish VRN1 and RTV1.

Thus a method involving use of PCR in obtaining nucleic acid according to the present invention may include:

(a) providing a preparation of plant nucleic acid, e.g. from a  
35 seed or other appropriate tissue or organ,

(b) providing a pair of nucleic acid molecule primers useful in (i.e. suitable for) PCR, at least one of said primers being a primer according to the present invention as discussed above,

- 5 (c) contacting nucleic acid in said preparation with said primers under conditions for performance of PCR,  
(d) performing PCR and determining the presence or absence of an amplified PCR product. The presence of an amplified PCR product may indicate identification of a variant.

10

In all cases above, if need be, clones or fragments identified in the search can be extended. For instance if it is suspected that they are incomplete, the original DNA source (e.g. a clone library, mRNA preparation etc.) can be revisited to  
15 isolate missing portions e.g. using sequences, probes or primers based on that portion which has already been obtained to identify other clones containing overlapping sequence.

If a putative naturally occurring homologous sequence is  
20 identified, its role in vernalisation can be confirmed, for instance by methods analogous to those used in the Examples below, or by generating mutants of the gene (e.g. by screening the available insertional-mutant collections) and analyzing these for their ability to respond to vernalization, possibly  
25 in the presence and absence of other alleles such as *vrn1*. Alternatively the role can be inferred from mapping *vrn* mutants to see if the homologue lies at or close to an appropriate locus.

30 In a further embodiment, antibodies raised to a VRN1 polypeptide or peptide can be used in the identification and/or isolation of variant polypeptides, and then their encoding genes. Thus, the present invention provides a method of identifying or isolating VRN1 or variant thereof,  
35 comprising screening candidate polypeptides with a polypeptide



comprising the antigen-binding domain of an antibody (for example whole antibody or a fragment thereof) which is able to bind said VRN1 polypeptide or variant thereof, or preferably has binding specificity for such a polypeptide. Methods of obtaining antibodies are described hereinafter.

Candidate polypeptides for screening may for instance be the products of an expression library created using nucleic acid derived from an plant of interest, or may be the product of a purification process from a natural source. A polypeptide found to bind the antibody may be isolated and then may be subject to amino acid sequencing. Any suitable technique may be used to sequence the polypeptide either wholly or partially (for instance a fragment of the polypeptide may be sequenced). Amino acid sequence information may be used in obtaining nucleic acid encoding the polypeptide, for instance by designing one or more oligonucleotides (e.g. a degenerate pool of oligonucleotides) for use as probes or primers in hybridization to candidate nucleic acid.

This aspect of the invention further includes an isolated nucleic acid comprising a sequence which is complementary to any of those isolated or obtained as above. The 'complement' in each case is the same length as the reference, but is 100% complementary thereto whereby by each nucleotide is base paired to its counterpart i.e. G to C, and A to T or U.

As used hereinafter, unless the context demands otherwise, the term "VRN1" is intended to cover any of the nucleic acids of the invention described above, including functional variants.

In one aspect of the present invention, the VRN1 nucleic acid described above is in the form of a recombinant and preferably replicable vector.

"Vector" is defined to include, inter alia, any plasmid, cosmid, phage or *Agrobacterium* binary vector in double or single stranded linear or circular form which may or may not be self transmissible or mobilizable, and which can transform a prokaryotic or eukaryotic host either by integration into the cellular genome or exist extrachromosomally (e.g. autonomous replicating plasmid with an origin of replication).

Generally speaking, those skilled in the art are well able to construct vectors and design protocols for recombinant gene expression. Suitable vectors can be chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, terminator fragments, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate. For further details see, for example, *Molecular Cloning: a Laboratory Manual*: 2nd edition, Sambrook et al, 1989, Cold Spring Harbor Laboratory Press or *Current Protocols in Molecular Biology*, Second Edition, Ausubel et al. eds., John Wiley & Sons, 1992.

Specifically included are shuttle vectors by which is meant a DNA vehicle capable, naturally or by design, of replication in two different host organisms, which may be selected from actinomycetes and related species, bacteria and eucaryotic (e.g. higher plant, mammalian, yeast or fungal cells).

A vector including nucleic acid according to the present invention need not include a promoter or other regulatory sequence, particularly if the vector is to be used to introduce the nucleic acid into cells for recombination into the genome.

Preferably the nucleic acid in the vector is under the control of, and operably linked to, an appropriate promoter or other regulatory elements for transcription in a host cell such as a

microbial, e.g. bacterial, or plant cell. The vector may be a bi-functional expression vector which functions in multiple hosts. In the case of genomic DNA, this may contain its own promoter or other regulatory elements and in the case of cDNA  
5 this may be under the control of an appropriate promoter or other regulatory elements for expression in the host cell

By "promoter" is meant a sequence of nucleotides from which transcription may be initiated of DNA operably linked  
10 downstream (i.e. in the 3' direction on the sense strand of double-stranded DNA).

"Operably linked" means joined as part of the same nucleic acid molecule, suitably positioned and oriented for  
15 transcription to be initiated from the promoter. DNA operably linked to a promoter is "under transcriptional initiation regulation" of the promoter.

In a preferred embodiment, the promoter is an inducible  
20 promoter.

The term "inducible" as applied to a promoter is well understood by those skilled in the art. In essence, expression under the control of an inducible promoter is "switched on" or  
25 increased in response to an applied stimulus. The nature of the stimulus varies between promoters. Some inducible promoters cause little or undetectable levels of expression (or no expression) in the absence of the appropriate stimulus. Other inducible promoters cause detectable constitutive  
30 expression in the absence of the stimulus. Whatever the level of expression is in the absence of the stimulus, expression from any inducible promoter is increased in the presence of the correct stimulus.

35 Thus this aspect of the invention provides a gene construct,

preferably a replicable vector, comprising a promoter (optionally inducible) operably linked to a nucleotide sequence provided by the present invention, such as the VRN1 gene or a variant thereof.

5

Particularly of interest in the present context are nucleic acid constructs which operate as plant vectors. Specific procedures and vectors previously used with wide success upon plants are described by Guerineau and Mullineaux (1993) (Plant transformation and expression vectors. In: Plant Molecular Biology Labfax (Croy RRD ed) Oxford, BIOS Scientific Publishers, pp 121-148). Suitable vectors may include plant viral-derived vectors (see e.g. EP-A-194809).

10

15 Suitable promoters which operate in plants include the Cauliflower Mosaic Virus 35S (CaMV 35S). Other examples are disclosed at pg 120 of Lindsey & Jones (1989) "Plant Biotechnology in Agriculture" Pub. OU Press, Milton Keynes, UK. The promoter may be selected to include one or more  
20 sequence motifs or elements conferring developmental and/or tissue-specific regulatory control of expression. Inducible plant promoters include the ethanol induced promoter of Caddick et al (1998) Nature Biotechnology 16: 177-180.

20

25 If desired, selectable genetic markers may be included in the construct, such as those that confer selectable phenotypes such as resistance to antibiotics or herbicides (e.g. kanamycin, hygromycin, phosphotricin, chlorsulfuron, methotrexate, gentamycin, spectinomycin, imidazolinones and  
30 glyphosate).

30

The present invention also provides methods comprising introduction of such a construct into a plant cell or a microbial cell and/or induction of expression of a construct  
35 within a plant cell, by application of a suitable stimulus

35

e.g. an effective exogenous inducer.

In a further aspect of the invention, there is disclosed a host cell containing a heterologous construct according to the present invention, especially a plant or a microbial cell.

The term "heterologous" is used broadly in this aspect to indicate that the gene/sequence of nucleotides in question (e.g. encoding VRN1) have been introduced into said cells of the plant or an ancestor thereof, using genetic engineering, i.e. by human intervention. A heterologous gene may replace an endogenous equivalent gene, i.e. one which normally performs the same or a similar function, or the inserted sequence may be additional to the endogenous gene or other sequence.

Nucleic acid heterologous to a plant cell may be non-naturally occurring in cells of that type, variety or species. Thus the heterologous nucleic acid may comprise a coding sequence of or derived from a particular type of plant cell or species or variety of plant, placed within the context of a plant cell of a different type or species or variety of plant. A further possibility is for a nucleic acid sequence to be placed within a cell in which it or a homologue is found naturally, but wherein the nucleic acid sequence is linked and/or adjacent to nucleic acid which does not occur naturally within the cell, or cells of that type or species or variety of plant, such as operably linked to one or more regulatory sequences, such as a promoter sequence, for control of expression.

The host cell (e.g. plant cell) is preferably transformed by the construct, which is to say that the construct becomes established within the cell, altering one or more of the cell's characteristics and hence phenotype e.g. with respect to a vernalisation response.

Nucleic acid can be introduced into plant cells using any

- suitable technology, such as a disarmed Ti-plasmid vector carried by *Agrobacterium* exploiting its natural gene transfer ability (EP-A-270355, EP-A-0116718, NAR 12(22) 8711 - 87215 1984), particle or microprojectile bombardment (US 5100792, 5 EP-A-444882, EP-A-434616) microinjection (WO 92/09696, WO 94/00583, EP 331083, EP 175966, Green et al. (1987) *Plant Tissue and Cell Culture*, Academic Press), electroporation (EP 290395, WO 8706614 Gelvin Debeyser) other forms of direct DNA uptake (DE 4005152, WO 9012096, US 4684611), liposome mediated 10 DNA uptake (e.g. Freeman et al. *Plant Cell Physiol.* 29: 1353 (1984)), or the vortexing method (e.g. Kindle, *PNAS U.S.A.* 87: 1228 (1990d) Physical methods for the transformation of plant cells are reviewed in Oard, 1991, *Biotech. Adv.* 9: 1-11.
- 15 *Agrobacterium* transformation is widely used by those skilled in the art to transform dicotyledonous species.

Recently, there has also been substantial progress towards the routine production of stable, fertile transgenic plants in 20 almost all economically relevant monocot plants (see e.g. Hiei et al. (1994) *The Plant Journal* 6, 271-282)). Microprojectile bombardment, electroporation and direct DNA uptake are preferred where *Agrobacterium* alone is inefficient or ineffective. Alternatively, a combination of different 25 techniques may be employed to enhance the efficiency of the transformation process, eg bombardment with *Agrobacterium* coated microparticles (EP-A-486234) or microprojectile bombardment to induce wounding followed by co-cultivation with *Agrobacterium* (EP-A-486233).

30 Preferred transformation protocols for brassicas, wheat, barley and rice may be found Becker et al., 1994 and references therein. However the skilled person will appreciate that the particular choice of a transformation 35 technology will be determined by its efficiency to transform

certain plant species as well as the experience and preference of the person practising the invention with a particular methodology of choice.

5 Thus a further aspect of the present invention provides a method of transforming a plant cell involving introduction of a construct as described above into a plant cell and causing or allowing recombination between the vector and the plant cell genome to introduce a nucleic acid according to the  
10 present invention into the genome.

The invention further encompasses a host cell transformed with nucleic acid or a vector according to the present invention (e.g comprising the VRN1 sequence) especially a plant or a  
15 microbial cell. In the transgenic plant cell (i.e. transgenic for the nucleic acid in question) the transgene may be on an extra-genomic vector or incorporated, preferably stably, into the genome. There may be more than one heterologous nucleotide sequence per haploid genome.

20

Generally speaking, following transformation, a plant may be regenerated, e.g. from single cells, callus tissue or leaf discs, as is standard in the art. Almost any plant can be entirely regenerated from cells, tissues and organs of the  
25 plant. Available techniques are reviewed in Vasil et al., *Cell Culture and Somatic Cell Genetics of Plants, Vol I, II and III, Laboratory Procedures and Their Applications*, Academic Press, 1984, and Weissbach and Weissbach, *Methods for Plant Molecular Biology*, Academic Press, 1989.

30

The generation of fertile transgenic plants has been achieved in the cereals rice, maize, wheat, oat, and barley (reviewed in Shimamoto, K. (1994) *Current Opinion in Biotechnology* 5, 158-162.; Vasil, et al. (1992) *Bio/Technology* 10, 667-674;  
35 Vain et al., 1995, *Biotechnology Advances* 13 (4): 653-671;

Vasil, 1996, *Nature Biotechnology* 14 page 702).

Plants which include a plant cell according to the invention are also provided. A plant according to the present invention  
5 may be one which does not breed true in one or more properties.

In addition to the regenerated plant, the present invention embraces all of the following: a clone of such a plant, seed,  
10 selfed or hybrid progeny and descendants (e.g. F1 and F2 descendants). The invention also provides a plant propagule from such plants, that is any part which may be used in reproduction or propagation, sexual or asexual, including cuttings, seed and so on. It also provides any part of these  
15 plants, which in all cases include the plant cell or heterologous VRN1 DNA described above.

Thus, one example of the above embodiment, would be to constitutively express the VRN1 protein in a transgenic plant  
20 e.g. by use of a fusion between the 35S promoter from cauliflower mosaic virus and the open reading frame from the VRN1 cDNA. Preferably this would use the binary vector SLJ1714 (Jones JDG, Shlummokov L, Carland F, English J, Scofield SR, Bishop GJ, Harrison K: Effective vectors for  
25 transformation, expression of heterologous genes, and assaying transposon excision in transgenic plants. *Transgenic Research* 1: 285-297 (1992)) using standard molecular techniques (Sambrook et al., 1989). In a further embodiment, inducible expression of the VRN1 protein is achieved using a gene  
30 fusions between the VRN1 open reading frame and the receptor domain of the rat glucocorticoid receptor (GR). An example of the use of this strategy to achieve inducible gene function can be found in Schena M, Lloyd AM, Davis RW: A steroid-inducible gene expression system for plant cells. *Proc Natl*  
35 *Acad Sci U S A* 88(23):10421-10425 (1991) and Simon R, Igeno



MI, Coupland G: Activation of floral meristem identity genes in Arabidopsis. Nature 384(6604): 59-62 (1996). Gene fusions can be tested, if desired, in the *vrn1-2* mutant allele of Arabidopsis by standard Agrobacterium mediate transfer

5 (Hoekema A, Hirsch PR, Hooykaas PJJ, Schilperroot A: A binary plant vector strategy based on separation of *vir*- and *T*-region of the Agrobacterium tumefaciens Ti-plasmid. Nature 303: 179-180 (1983)). The vernalization requirement of the different transgenic plants obtained will be analyzed compared to  
10 control (non-transformed) plants.

A further aspect of the present invention provides a method for assessing the vernalisation responsiveness of a plant, the method comprising the step of determining the presence and/or  
15 identity of a *VRN1* allele therein comprising the use of a nucleic acid as described above. Such a diagnostic test may be used with transgenic or wild-type plants. The use of diagnostic tests for alleles allows the researcher or plant breeder to establish, with full confidence and independent  
20 from time consuming biochemical tests, whether or not a desired allele is present in the plant of interest (or a cell thereof), whether the plant is a representative of a collection of other genetically identical plants (e.g. an inbred variety or cultivar) or one individual in a sample of  
25 related (e.g. breeders' selection) or unrelated plants.

The method may form part of a plant breeding scheme based on selection and selfing of desirable individuals. Reliable selection for appropriate *VRN1* alleles can be made in early  
30 generations and on more material than would otherwise be possible. This gain in reliability of selection plus the time saving by being able to test material earlier and without costly phenotype screening is of considerable value in plant breeding.

Nucleic acid-based determination of the presence or absence of one or more desirable alleles may be combined with determination of the genotype of the flanking linked genomic DNA and other unlinked genomic DNA using established sets of markers such as RFLPs, microsatellites or SSRs, AFLPs, RAPDs etc. This enables the researcher or plant breeder to select for not only the presence of the desirable allele but also for individual plant or families of plants which have the most desirable combinations of linked and unlinked genetic background. Such recombinations of desirable material may occur only rarely within a given segregating breeding population or backcross progeny. Direct assay of the locus as afforded by the present invention allows the researcher to make a step-wise approach to fixing (making homozygous) the desired combination of flanking markers and alleles, by first identifying individuals fixed for one flanking marker and then identifying progeny fixed on the other side of the locus all the time knowing with confidence that the desirable allele is still present.

The present disclosure provides sufficient information for a person skilled in the art to obtain genomic DNA sequence for any given new or existing allele and devise a suitable nucleic acid- and/or polypeptide-based diagnostic assay. In designing a nucleic acid assay account is taken of the distinctive variation in sequence that characterizes the particular variant allele (see e.g. Fig 7 and the allelic variations described therein).

The invention further provides a method of influencing or affecting the vernalisation response in a plant, the method including causing or allowing expression of a heterologous VRN1 nucleic acid sequence as discussed above within the cells of the plant. The method may include the use of VRN1 nucleic acid in conjunction with other genes affecting vernalisation

(e.g. VRN2). As discussed in the Examples below, VRN1 and VRN2 may act in separate and partially redundant vernalization-promoting pathways.

- 5 The step may be preceded by the earlier step of introduction of the VRN1 nucleic acid into a cell of the plant or an ancestor thereof. In addition to use of the nucleic acids of the present invention for production of functional VRN1 polypeptides (thereby enhancing the vernalisation response),  
10 the information disclosed herein may also be used to reduce the activity VRN1 activity in cells in which it is desired to do so (thereby inhibiting or destroying the vernalisation response).
- 15 For instance down-regulation of expression of a target gene may be achieved using anti-sense technology.

In using anti-sense genes or partial gene sequences to down-regulate gene expression, a nucleotide sequence is placed  
20 under the control of a promoter in a "reverse orientation" such that transcription yields RNA which is complementary to normal mRNA transcribed from the "sense" strand of the target gene. See, for example, Rothstein et al, 1987; Smith et al, (1988) *Nature* 334, 724-726; Zhang et al, (1992) *The Plant*  
25 *Cell* 4, 1575-1588, English et al., (1996) *The Plant Cell* 8, 179-188. Antisense technology is also reviewed in Bourque, (1995), *Plant Science* 105, 125-149, and Flavell, (1994) *PNAS USA* 91, 3490-3496.

- 30 An alternative to anti-sense is to use a copy of all or part of the target gene inserted in sense, that is the same, orientation as the target gene, to achieve reduction in expression of the target gene by co-suppression. See, for example, van der Krol et al., (1990) *The Plant Cell* 2, 291-  
35 299; Napoli et al., (1990) *The Plant Cell* 2, 279-289; Zhang et

al., (1992) *The Plant Cell* 4, 1575-1588, and US-A-5,231,020. Further refinements of the gene silencing or co-suppression technology may be found in W095/34668 (Biosource); Angell & Baulcombe (1997) *The EMBO Journal* 16,12:3675-3684; and Voinnet  
5 & Baulcombe (1997) *Nature* 389: pg 553.

Further options for down regulation of gene expression include the use of ribozymes, e.g. hammerhead ribozymes, which can catalyse the site-specific cleavage of RNA, such as mRNA (see  
10 e.g. Jaeger (1997) "The new world of ribozymes" *Curr Opin Struct Biol* 7:324-335, or Gibson & Shillitoe (1997) "Ribozymes: their functions and strategies form their use" *Mol Biotechnol* 7: 242-251.)

15 The complete sequence corresponding to the coding sequence (in reverse orientation for anti-sense) need not be used. For example fragments of sufficient length may be used. It is a routine matter for the person skilled in the art to screen fragments of various sizes and from various parts of the  
20 coding sequence to optimise the level of anti-sense inhibition. It may be advantageous to include the initiating methionine ATG codon, and perhaps one or more nucleotides upstream of the initiating codon. A further possibility is to target a conserved sequence of a gene, e.g. a sequence that is  
25 characteristic of one or more genes, such as a regulatory sequence.

The sequence employed may be about 500 nucleotides or less, possibly about 400 nucleotides, about 300 nucleotides, about  
30 200 nucleotides, or about 100 nucleotides. It may be possible to use oligonucleotides of much shorter lengths, 14-23 nucleotides, although longer fragments, and generally even longer than about 500 nucleotides are preferable where possible, such as longer than about 600 nucleotides, than

about 700 nucleotides, than about 800 nucleotides, than about 1000 nucleotides or more.

It may be preferable that there is complete sequence identity in the sequence used for down-regulation of expression of a target sequence, and the target sequence, although total complementarity or similarity of sequence is not essential. One or more nucleotides may differ in the sequence used from the target gene. Thus, a sequence employed in a down-regulation of gene expression in accordance with the present invention may be a wild-type sequence (e.g. gene) selected from those available, or a variant of such a sequence in the terms described above. The sequence need not include an open reading frame or specify an RNA that would be translatable.

Thus the present invention further provides the use of a variant VRN1 nucleotide sequence, or its complement, for down-regulation of gene expression, particularly down-regulation of expression of the VRN1 gene or homologue thereof, preferably in order to inhibit or suppress the vernalisation response in a plant.

Anti-sense or sense regulation may itself be regulated by employing an inducible promoter in an appropriate construct.

The present invention also encompasses the expression product of any of the coding VRN1 nucleic acid sequences disclosed and methods of making the expression product by expression from encoding nucleic acid therefore under suitable conditions, which may be in suitable host cells.

As described in the Examples, several features of VRN1 suggest that it is likely to serve as a modulator of transcription (e.g., as a "co-activator" or "co-repressor"), or in the least as a DNA-binding protein. These features include the presence

of the B3 domains; the homology of a portion of region 2 with c-myc, a transcription factor; the presence of a putative NLS, and the presence of putative signals for rapid protein degradation, which are common in transcription factors and other proteins of regulatory function (Chevaillier, 1993; Vierstra, 1996; Barnes and Gomes, 1995; Rechsteiner and Rogers, 1996; Gomes and Barnes, 1997).

The present invention also provides for the production and use of fragments of the full-length polypeptides disclosed herein, especially active portions thereof. An "active portion" of a polypeptide means a peptide which is less than said full length polypeptide, but which retains an essential biological activity. In particular, the active portion retains the ability to alter vernalization response in a plant, such as *Arabidopsis thaliana*.

A "fragment" of a polypeptide means a stretch of amino acid residues of at least about five to seven contiguous amino acids, often at least about seven to nine contiguous amino acids, typically at least about nine to 13 contiguous amino acids and, most preferably, at least about 20 to 30 or more contiguous amino acids.

Use of recombinant VRN1 protein, or a fragment (e.g the domains discussed above) thereof, as a DNA-binding protein, or more specifically a modulator of transcription, forms one aspect of the invention.

Fragments of the polypeptides may include one or more epitopes useful for raising antibodies to a portion of any of the amino acid sequences disclosed herein. Preferred epitopes are those to which antibodies are able to bind specifically, which may be taken to be binding a polypeptide or fragment thereof of

the invention with an affinity which is at least about 1000x that of other polypeptides.

Thus purified VRN1 protein, or a fragment or other variant thereof, e.g. produced recombinantly by expression from encoding nucleic acid therefor, may be used to raise antibodies employing techniques which are standard in the art. Antibodies and other polypeptides comprising antigen-binding fragments of antibodies may be used in identifying homologues from other plant species as discussed above.

Methods of producing antibodies include immunising a mammal (e.g. mouse, rat, rabbit, horse, goat, sheep or monkey) with the protein or a fragment thereof. Antibodies may be obtained from immunised animals using any of a variety of techniques known in the art, and might be screened, preferably using binding of antibody to antigen of interest.

For instance, Western blotting techniques or immunoprecipitation may be used (Armitage et al, 1992, Nature 357: 80-82). Antibodies may be polyclonal or monoclonal.

Antibodies may be modified in a number of ways. Indeed the term "antibody" should be construed as covering any specific binding substance having a binding domain with the required specificity. Thus, this term covers antibody fragments, derivatives, functional equivalents and homologues of antibodies, including any polypeptide comprising an immunoglobulin binding domain, whether natural or synthetic.

As an alternative or supplement to immunising a mammal, antibodies with appropriate binding specificity may be obtained from a recombinantly produced library of expressed immunoglobulin variable domains, e.g. using lambda bacteriophage or filamentous bacteriophage which display

functional immunoglobulin binding domains on their surfaces;  
for instance see WO92/01047.

Specific binding members such as antibodies and polypeptides

5 comprising antigen binding domains of antibodies that bind and  
are preferably specific for a VRN1 polypeptide or variant  
thereof represent further aspects of the present invention, as  
do their use and methods which employ them.

10 The above description has generally been concerned with the  
coding parts of the VRN1 gene and variants and products  
thereof. Also embraced within the present invention are  
untranscribed parts of the gene.

15 Thus a further aspect of the invention is a nucleic acid  
molecule encoding the promoter of the VRN1 gene, which is  
believed to be present in the sequence shown in Annex I  
(which begins at the end of the *LARS1* gene).

20 As described in the Examples below, The VRN1 promoter region  
and VRN1 intron 1 were found to contain a variety of potential  
binding sites including low temperature response elements;  
binding sites for the *Arabidopsis* dehydration- and ABA-  
responsive gene rd22; one binding site for *Arabidopsis* Myb2, a  
25 transcription factor involved in regulation of genes  
responsive to water stress; H-box and TCA-1 binding sites  
(that may be induced by wounding and abiotic stress); and  
ICE-boxes (a consensus promoter element found in several cold-  
inducible genes).

30

These control elements are likely to dictate the conditions in  
which expression of the VRN1 transcript is obtained. For  
example, VRN1 may perhaps be induced by cold and/or drought  
treatment, or simply by application of ABA, and use of the  
35 promoter or a part thereof for induction of transcription



under any of these conditions forms one aspect of the present invention

Analysis of the upstream region will reveal control regions  
5 for gene expression including control regions common to many  
genes (i.e TATA and CAAT boxes) and other control regions,  
usually located from 1 to 10,000, such as 1 to 1000 or 50 to  
500 nucleotides upstream of the start of transcription. To  
find minimal elements or motifs responsible for regulation,  
10 restriction enzyme or nucleases may be used to digest a  
nucleic acid molecule, or mutagenesis may be employed,  
followed by an appropriate assay (for example using a reporter  
gene such as luciferase) to determine promoter activity. The  
control region may also be mutated to identify specific  
15 subregions responsible for transcriptional control. This may  
be achieved by a number of techniques well known in the art as  
such, including DNase protection footprint assays, in which  
the control region is brought into contact with an extract  
from a cell in which the VRN1 gene is actively expressed, and  
20 the regions of the control region which bind factors in that  
extract is determined.

Nucleic acid comprising these elements or motifs forms one  
part of the present invention.

25

"Promoter activity" is used to refer to ability to initiate  
transcription under appropriate conditions e.g. optionally in  
the presence of an inducer. The level of promoter activity is  
quantifiable for instance by assessment of the amount of mRNA  
30 produced by transcription from the promoter or by assessment  
of the amount of protein product produced by translation of  
mRNA produced by transcription from the promoter. The amount  
of a specific mRNA present in an expression system may be  
determined for example using specific oligonucleotides which  
35 are able to hybridise with the mRNA and which are labelled or

may be used in a specific amplification reaction such as the polymerase chain reaction.

Those skilled in the art are well aware of a multitude of possible reporter genes and assay techniques which may be used to determine promoter activity. Any suitable reporter/assay may be used and it should be appreciated that no particular choice is essential to or a limitation of the present invention. Also provided is a nucleic acid construct, preferably an expression vector, including the VRN1 promoter (or active fragment or variant thereof able to promote transcription) operably linked to a heterologous gene, e.g. a coding sequence, which is preferably not the coding sequence with which the promoter is operably linked in nature.

The invention will now be further described with reference to the following non-limiting Figures and Examples. Other embodiments of the invention will occur to those skilled in the art in the light of these.

#### FIGURES & SEQUENCE ANNEXES

Fig 1: Vernalization phenotype of *vrn1* mutant under LDs and SDs; vernalization phenotype of *vrn1-1* allele compared to *vrn1-2* allele.

Fig 2: Genetic map of the position of VRN1 on chromosome III in relation to markers used for mapping. The markers (shown on right) were scored on a population of 494 F2 plants from a cross between *vrn1-1 fac1* x *fca-10*. The distance in cM between each marker is shown on the left.

Fig 3: Physical map of the region containing VRN1.

Fig 4: Complementation of the *vrn1-1* mutant phenotype by cosmids 8H8 and 10F10. Following vernalization, *fca-1* plants flower early and *vrn1-1 fca-1* plants flower late.

Representative T2 lines in which cosmid 8H8 or 10F10 has been transformed in *vrn1-1 fca-1* plants show the expected ratio (approx. 3:1) of early-to-late flowering plants.

Fig 5: Sequenced region, and predicted ORFs in the vicinity of *VRN1*. Overlap between cosmids was initially determined by XbaI + XhoI digestion and Southern blotting. Sequencing of cosmid DNA confirmed these results and revealed the complementing region as 6565 bp. ORF1 was subsequently shown to be *VRN1*.

Fig 6: Structure of the *VRN1* gene and transcript, and positions of the *vrn1-1* and *vrn1-2* mutations.

Fig 7: The putative *VRN1* transcript and its deduced amino acid sequence.

Fig 8: Alignment of *VRN1* and *RTV1*

Annex I: this shows contig 29 [bp 1501-6500]) derived from Ler *VRN1* genomic DNA. The *VRN1* promoter is present in the region between about nucleotides 1 to 1879.

## EXAMPLES

### Example 1 - Isolation of the *vrn1* mutants

The *vrn1* mutation was selected from mutagenized populations of *Arabidopsis thaliana* (L.) Heynh (Landsberg erecta ecotype) plants on the basis of its impairment of the acceleration of flowering following a six week cold treatment (vernalization).

Mutants were subsequently analyzed for flowering time in the absence of vernalization in order to confirm that the induced defect was specific to the process of vernalization and not due to a general late-flowering mutation (Chandler et al., 1996).

Two recessive alleles of *vrn1* have been identified: (1) *vrn1-1* was isolated by mutagenising *fca-1* seeds with EMS, as described (Chandler et al., 1996), and (2) *vrn1-2* was isolated by mutagenising *fca-1* seeds with gamma irradiation. The *vrn1-1 fca-1* line used here was backcrossed to *fca-1* two times prior to genetic mapping. Subsequently, *vrn1-1 fca-1* was further backcrossed to *fca-1* (six times in total) and *vrn1-2 fca-1* was backcrossed two times in total.

#### Example 2 - Characterization of the *vrn1* phenotype

The vernalization dose-response phenotype of *vrn1* mutant plants was investigated by examining their flowering time in response to differing durations of vernalization treatment. Flowering time was measured in two ways: (1) as the total number of vegetative leaves produced prior to flowering (LN), and (2) as the time in days from the end of the vernalization treatment to the appearance of the first floral bud (FT). In all experiments these two measures were positively correlated, so only LN is given in order to more easily facilitate comparison between experiments.

Two types of experiment were conducted: (1) a dose-response analysis of *vrn1-1 fca-1* and *vrn1-2 fca-1* examined under long day (LD) growth conditions (Figure 1A), and (2) the effect of 6 weeks of vernalization on *vrn1-1* in the absence of *fca-1* examined under short day (SD) growth conditions (Figure 1B). In the LD experiment shown in Figure 1A, with no

vernalization (0 weeks), both *vrn1-1 fca-1* and *vrn1-2 fca-1* mutant plants flowered very slightly earlier compared to the parental *fca-1* controls, although in other experiments *vrn1-1 fca-1* and *vrn1-2 fca-1* mutant plants flowered at approximately the same time as *fca-1* with no vernalization. In contrast, following vernalization, *fca-1* plants showed a marked reduction in leaf number ( $\approx 66\%$  after 6 weeks of vernalization), while *vrn1-1 fca-1* and *vrn1-2 fca-1* mutant plants showed a much reduced response ( $\approx 14\%$  and  $\approx 27\%$  after 6 weeks of vernalization, respectively). Therefore, both alleles of *vrn1* are dramatically impaired in their response to vernalization, with *vrn1-1* being more severe than *vrn1-2*.

In the SD experiment shown in Figure 1B, the wild type Ler plants exhibited a  $\approx 49\%$  reduction in leaf number after a vernalization treatment of six weeks compared to unvernallized plants. However, *vrn1-1* mutant plants showed only a  $\approx 18\%$  reduction under the same conditions. In addition, this experiment shows that the phenotype of *vrn1-1* does not depend on the presence of the *fca-1* mutation or on long day photoperiods. *vrn1-1* was also combined with other late flowering, vernalization-responsive mutations (*fve-1*, *ld-3*, *fwa-1*, *fe-1*, *fpa-2*, and *ft-1*) and was shown to impair the vernalization response of these mutants as well (Chandler et al., 1996).

### Example 3 - Genetic mapping of VRN1

The *VRN1* gene was initially mapped to the top arm of chromosome III, between RLFP markers mi207 and mi339, using a relatively small F2 population (77 plants) derived from a cross between *vrn1-1 fca-1* and *fca-10*, as described (Chandler et al., 1996). A larger population (494 F2 plants) derived from the same cross was then used to finely map the position

of *VRN1* (Figure 2). The dearth of available genetic markers in this region necessitated the development of several new genetic markers that were polymorphic between the Ler and Ws ecotypes (Table 1). As a first step, two markers flanking  
5 *VRN1*, *ATHCHIB* (SSLP) and *g4711* (CAPS) were used to screen the population for recombinants in this  $\approx 18$  cM interval. Approximately 170 recombinant chromosomes were identified. Next, the markers indicated in Figure 2 were used with these recombinants to define the position of *VRN1* to the  $\approx 0.5$  cM  
10 interval between *mi339* (2 recombinants to the north) and *pKS1240* (one recombinant to the south). The CAPS marker *agp14*, corresponding to a dioxygenase gene, was genetically inseparable from *VRN1* (Figure 2).

#### 15 Example 4 - Physical mapping of *VRN1*

The interval between *mi339* and *pKS1240* fell in a gap between Contig 3 and Contig 4 of the CIC YAC coverage of chromosome III (Camilleri et al., 1998) so therefore no physical map data  
20 was available. Initially, an attempt was made to fill the gap using YAC clones other than those derived from the CIC library (i.e., yUP, EW, and EG), but this genomic region was apparently not represented in any of these libraries. Therefore, a physical map of the interval was constructed  
25 using IGF (Mozo et al., 1998) and TAMU (website address: [genome-www.stanford.edu/Arabidopsis/ww/Vol2/choi.html](http://genome-www.stanford.edu/Arabidopsis/ww/Vol2/choi.html)) BAC clones. Marker *mi339* was used to screen the BAC libraries and to initiate a walk towards *pKS1240*. BAC contigs (Figure 3) were assembled by hybridizing BACs to end probes developed by  
30 iPCR (Table 2) and by using publicly available BAC end sequence data (from TIGR; website address [www.tigr.org/tdb/at/atgenome/bac\\_end\\_search/bac\\_end\\_search.htm](http://www.tigr.org/tdb/at/atgenome/bac_end_search/bac_end_search.htm) 1) as the basis for designing oligonucleotide primers for PCR (Table 2). The size of individual BAC clones was determined by  
35 hexagonal pulsed-field gel electrophoresis (Maule, 1997). The

≈0.5 cM interval between mi339 and pKS1240 containing the *VRN1* gene was therefore found to correspond to ≈120 kb of genomic DNA.

5 In preparation for cosmid complementation experiments, a Ler genomic library in the cosmid 04541 binary vector (Macknight et al., 1997) was initially screened using the following probes: BAC T24F13, mi339, agp14, and pKS1240. Putative positive clones were verified on Southern blots and the  
10 overlap between individual cosmids determined by either hybridization with DNA probes or with PCR primers designed from BAC- and cosmid- end sequence data (Table 2). The insert sizes of individual cosmid clones was determined by digestion with XbaI + XhoI followed by standard agarose gel  
15 electrophoresis using lambda DNA cut with HindIII as a standard. A complete cosmid contig was generated over the ≈120 kb region (Figure 3).

Example 5 - Cosmid complementation of the *vrn1* phenotype

20

Eight cosmids (39K3, 8H8, 10F10, 42A10, 2P5, 19D3, 27J7, 67N6) centered around the marker agp14 were transformed into *vrn1-1 fca-1* plants by *Agrobacterium tumefaciens*-infection of root tissue (Hooykaas, 1989). In order to test if any of these  
25 cosmids rescued the mutant phenotype of *vrn1-1 fca-1*, T2 seed (from individual T1 kanamycin resistant transformants) was sown on soil and vernalized for 5 weeks. Seedlings were then transferred to LD conditions, and pricked out into individual compartments of divided trays after about a week of growth.

30

The total leaf number prior to flowering was determined, and cosmids were scored as complementing if the segregation ratio of early to late plants (compared to *fca-1* and *vrn1-1 fca-1* controls) was approximately 3:1 or greater. Eight independent lines containing cosmid 8H8, eight independent lines

35

containing cosmid 10F10, and three independent lines

containing cosmid 39K3 were found to rescue mutant phenotype of *vrn1-1 fca-1*. Lines containing the other five cosmids did not complement the *vrn1-1* phenotype (Figure 3). Analysis of the flowering time segregation in typical 8H8 and 10F10

5 complementing lines is shown in Figure 4. The presence of each cosmid in complementing lines (T2 plants) was confirmed by a cosmid-specific diagnostic

PCR, comprising an insert specific primer 8H8DIAG1

10 (ACCTGCTTCTGCCAACCGCTC) and 10F10DIAG1

(AGTTCGCTCTTGCTGTTTTTTTCCC) (corresponding to a portion of the Ler genomic DNA) and a primer BACT 7U (CCTCTTCGCTATTACGCCAG) present in the cosmid vector (see "cosmid complementation" under "materials and methods" below).

15 Example 6 - Analysis of genomic DNA corresponding to the complementing region

(a) Sequencing of cosmid DNA

20 The region of chromosome III corresponding to the cosmid contig surrounding *VRN1* (Figure 3) had apparently not previously been sequenced. Therefore the insert DNA from cosmids 8H8, 10F10, and 39K3 (derived from Ler genomic DNA) was sequenced by a combination of primer walking and shotgun strategies (Table 3), resulting in three contigs of sequence (Figure 5). The total amount of new *Arabidopsis* genomic sequence obtained was 20950 bp.

30 (B) Identification of candidate ORFs in genomic sequence

As new genomic sequence data was obtained it was analyzed in several ways in order to identify potential open reading frames (ORFs) and genes. Firstly, homology searches were carried out using the BLAST and FASTA computer programs



available from the *Arabidopsis thaliana* Database (AtDB; website address: genome-  
www.stanford.edu/Arabidopsis/seqtools.html) and National Center for Biotechnology Information (NCBI; website address:  
5 www.ncbi.nlm.nih.gov/BLAST/). Using these programs, genomic sequence in the *VRN1* region was compared (1) the *Arabidopsis* EST database, and (2) the database of all non-redundant Genbank sequences. Secondly, searches were carried out using the NetPlantGene website address:

10 www.cbs.dtu.dk/NetPlantGene.html), BCM Gene Finder website address: (dot.imgen.bcm.tmc.edu:9331/gene-finder/gf.html), and GENESCAN website address: genomic.stanford.edu/GENSCANW.html) computer programs which are designed to recognize features of eukaryotic genes, such as intron-exon boundaries, ORFs and  
15 polyadenylation signals. The results of these analyses are summarized in Figure 5 ("Predicted ORFs"). The sequenced region (contigs 29, 2, and 4) was found to contain  $\approx 8$  potential genes. Three of these, *agp14*, *LARS1*, and *ORF1* (later identified as *VRN1*) were represented by ESTs in the GenBank  
20 EST database.

Example 7 - Identification of the *vrn1-1* and *vrn1-2* mutations and determination of the *VRN1* gene structure

25 (A) *Finding mutations in vrn1 mutant plants*

The three cosmids which rescued the *vrn1-1 fca-1* mutant phenotype (8H8, 10F10, 39K3) were subjected to restriction analysis using XbaI and XhoI (Figure 5) and the region of  
30 overlap between these three cosmids found to be  $\approx 6.5$  kb. The ORF analysis indicated that this 6.5 kb interval contained the 3' end of the *LARS1* gene (a dioxygenase closely related to *agp14*), the 3' end of a hypothetical gene of unknown function, and the entire structure of another gene, "*ORF1*" (Figure 5).

In order to determine if either *LARS1* or *ORF1* corresponded to *VRN1*, a search for the presence of mutations in these genes in *vrn1-1 fca-1* and *vrn1-2 fca-1* mutant plants was carried out. PCR primers initially used in the sequencing of cosmid DNA (Table 3) were now used to amplify products from *vrn1-1 fca-1* and *vrn1-2 fca-1* genomic DNA. Overlapping products that encompassed the entire predicted ORF of *LARS1* and *ORF1* were sequenced on both strands and compared to the Ler-derived cosmid sequence for the presence of differences corresponding to mutations. No mutations were found in the *LARS1* gene, but in *ORF1*, a 1bp nonsense mutation was found in *vrn1-1 fca-1*-derived DNA and a 1bp deletion was found in *vrn1-2 fca-1*-derived DNA (Figure 6). Each of these putative mutations were then confirmed by sequencing four more independent PCR products on both strands. The effect of the *vrn1-1* and *vrn1-2* mutations on the encoded VRN1 protein is described in Example 8.

(B) Determining the structure of the VRN1 gene

The structure of the *VRN1* gene and putative transcript was determined by a combination of (1) RT-PCR analysis, (2) 3'-RACE analysis, and (3) analysis of partial cDNA clones represented in the GenBank *Arabidopsis* EST database (Table 4). These techniques revealed the sequence of the *VRN1* transcript and by comparing this sequence with the *VRN1* genomic sequence, the intron/exon boundaries were determined (Figure 6, Figure 7). The results obtained by these approaches were all in agreement, i.e., the intron-exon boundaries and point of polyadenylation determined by RT-PCR and 3'-RACE were identical to those determined through analysis of EST clones corresponding to *VRN1* cDNA, although the 5' transcription start site of the *VRN1* gene was not definitively determined by the experiments. Within the putative *VRN1* transcript (Figure

7), the Ler-derived sequence obtained by RT-PCR and the Columbia-derived sequence obtained by the sequencing of EST clones were 100% identical.

5 The *VRN1* gene is comprised of 5 exons and covers  $\approx 3.0$  kb of genomic DNA from the putative transcription start to the point of polyadenylation (see Annex I). Introns 2, 3, and 4 are a typical size for an *Arabidopsis* gene ( $\approx 100$  bp), while intron 1 is quite large:  $\approx 1.2$  kb (Figure 6). The 5- and 3-UTR of the  
10 *VRN1* transcript are also somewhat larger than average:  $\approx 270$  and  $\approx 200$  bp, respectively (Figures 6, 7).

The *VRN1* promoter region (from end of *LARS1* gene to *VRN1* translation initiation codon) and *VRN1* intron 1 were analyzed  
15 for binding sites of known plant transcription factors and known promoter elements using the Web Signal Scan program and PLACE database website address:  
([www.dna.affrc.go.jp/htdocs/PLACE/signalscan.html](http://www.dna.affrc.go.jp/htdocs/PLACE/signalscan.html)). These regions of *VRN1*, which may potentially specify the expression  
20 of the *VRN1* gene, were found to contain the following potential binding sites: (1) two low temperature response elements (LTREs; also known as CRT/DREs), found in several cold-induced genes from *Arabidopsis*, *Brassica napus*, and barley and bound by the transcription factor CBF1 (Baker et al., 1994; Stockinger et al., 1997; Jiang et al., 1996; Nordin et al., 1993), (2) three binding sites for the *Arabidopsis* dehydration- and ABA-responsive gene *rd22* (Abe et al., 1997), (3) one binding site for *Arabidopsis* Myb2, a transcription factor involved in regulation of genes responsive to water  
25 stress (Urao et al., 1993), (4) one H-box and three TCA-1 binding sites, promoter elements found in several tobacco, barley, and bean (*P. vulgaris*) genes that are induced by wounding and abiotic stress (Loake et al., 1992; Mhiri et al., 1997; Goldsbrough et al., 1993), and (5) three ICE-boxes, a  
30

consensus promoter element found in several cold-inducible genes (G.J. Warren, unpublished). It is interesting that control elements for both cold- and drought-inducible genes are present within the *VRN1* promoter and intron 1, as these conditions are known to induce several genes involved in acclimation to freezing temperatures (Thomashow, 1994), and ABA signaling is involved (Gilmour and Thomashow, 1991). These control elements are likely to dictate the conditions in which expression of the *VRN1* transcript is obtained. For example, *VRN1* may perhaps be induced by cold and/or drought treatment, or simply by application of ABA.

Taken together, the presence of mutations within *ORF1* (the only predicted gene that was completely contained within the complementing region) in genomic DNA derived from *vrn1-1* and *vrn1-2* mutant plants confirmed that *ORF1* corresponds to the *VRN1* gene.

This may be readily confirmed by introduction of the ORF (in sense and antisense) into *Arabidopsis* (cf. Example 5 above). Constructs may be based on the pGreen0029 vector which drives expression of the cloned in gene with a double 35S promoter and terminator derived from CaMV. This vector, and how to obtain it, is discussed in detail in WO 99/27120 (Plant Bioscience Limited).

(1) Genomic sense construct: the unspliced (genomic) *VRN1* ORF in the sense orientation in order to produce high levels of functional *VRN1* product. This construct will be put into *vrn1-1 fca-1* and *vrn1-2 fca-1* plants.

(2) cDNA sense construct: the spliced (cDNA) *VRN1* ORF in the sense orientation in order to produce high levels of

functional VRN1 product. This construct will be put into *vrn1-1 fca-1* and *vrn1-2 fca-1* plants.

- 5 (3) cDNA antisense construct: the spliced (cDNA) VRN1 ORF in the antisense orientation in order to repress the normal expression of VRN1 and diminish the amount of functional VRN1 product. This construct will be put into *fca-1* and Ler plants.

10 As an alternative to a constitutive promoter, it may be desirable to use an inducible promoter, such as one which is controlled by application of the molecule dexamethasone.

Example 8 - Analysis of the putative protein encoded by the VRN1 gene

15

The deduced amino acid sequence of VRN1 (Figure 7) was compared with the entire GenBank database (NCBI) using the BLASTP and TBLASTN programs.

- 20 (A) VRN1: domain structure, sequence features, and similarity to other known and hypothetical sequences

The VRN1 gene encodes a putative protein of 341aa (calculated MW=39278 Da) that is basic (pI=9.1), and is comprised of at least three regions. Region 1 (residues 2-94 in Figure 7) and 3 (residues 239-332) which are homologous to each other and are related to the B3 DNA-binding domain originally found in the maize transcription factor VIVIPAROUS1 (VP1; McCarty et al., 1991). Domains similar to the B3 domain of VP1 have subsequently been found in several *Arabidopsis* transcription factors or putative transcription factors such as ABI3 (the *Arabidopsis* orthologue of maize VP1, (Giraudat et al., 1992), auxin response factors (Ulmasov et al., 1997), IAA response factors (Kim et al., 1997; Abel et al., 1994; Guilfoyle et

30

al., 1998), FUSCA3 (Luerksen et al., 1998), and RAVs (Kagaya et al., 1999). Several of these proteins have been shown to bind DNA in sequence-specific manner via their B3 domain (e.g., Kagaya et al., 1999; Suzuki et al., 1997; Ulmasov et al., 1997).

The B3 DNA-binding domain appears to be specific to plants (Suzuki et al., 1997), and analysis of translated nucleotide sequences (i.e., hypothetical proteins) in the GenBank databases has revealed at least 22 *Arabidopsis* sequences that contain B3 domains, as well as EST sequences from several other plant species such as *Brassica oleracea*, hybrid aspen (*Populus tremula* x *P. tremuloides*), and tomato. While VRN1 contains two B3 domains, most characterized and hypothetical amino acid sequences were found to contain only one B3 domain, and some were found to contain more than two. The B3 domain appears to be "defined" by a number of conserved positions (results not shown) rather than sequence identity over the whole domain. Therefore, BLAST scores between the sequences shown tested are only marginally significant (on the order of  $10^{-6}$  to  $10^{-1}$ ). The C-terminal portion of the B3 domain is more conserved than the N-terminal portion.

Phylogenetic analysis of B3 domains (results not shown) using the Clustal method suggests that *Arabidopsis* B3-containing proteins fall into several groups: (1) ABI3- and FUSCA3-like B3s, (2) auxin response factor- (ARFs) and IAA-inducible protein-like B3s, (3) RAV1-like B3s, and (4) at least four uncharacterized groups, which include the VRN1-like B3s. It is likely that through evolution the B3 domain has been recruited in different ways by proteins involved in diverse plant processes.

Region 2 of VRN1 (residues 95-238), which lies between the two B3 domains (Figure 7), is not obviously related to any domain

of known function, nor does it have obvious features of a transcriptional activation or repression domain. Nonetheless, region 2 does contain several sequence features and motifs of interest, including a putative nuclear localization signal (NSL), two putative PEST regions (identified using the PEST Sequence Utility (website address: [www.lif.icnet.uk/LRITu/projects/pest/](http://www.lif.icnet.uk/LRITu/projects/pest/)) based on Rechsteiner and Rogers, 1996; Rogers et al., 1986), and three RXXL motifs also associated with rapid protein degradation (Cooper et al., 1997) (Figure 7). Interestingly, the second PEST region of VRN1 contains a potential protein kinase C (PKC) phosphorylation site (residues 176-178 in Figure 7). There are several examples in the literature for regulation of the cellular "lifespan" of proteins by phosphorylation of PEST regions (e.g., McKinsey et al., 1997; Koepp et al., 1999; Yaglom et al., 1996; Marchal et al., 1998; Liu et al., 1997). For example, in the case of I $\kappa$ B, stimulation of cell surface receptors by cytokines initiates a signal transduction cascade that phosphorylates I $\kappa$ B at two specific serine residues in the PEST region, triggering the polyubiquitination of nearby lysine residues and ultimately proteolysis (McKinsey et al., 1997; Laney and Hochstrasser, 1999).

Analysis of the physiochemical characteristics of VRN1 suggest that the two B3 domains are basic (average pI $\approx$ 9.5) and slightly hydrophobic in character, while region 2 is slightly acidic (pI $\approx$ 6.3) and somewhat hydrophilic and therefore likely to be on the surface of the molecule and exposed to the aqueous environment. Interestingly, unlike the B3 domains which appear to be specific to plants, BLAST searches against GenBank (NCBI) with region 2 of VRN1 picked up no significant hits from plants (except for RTV1, see below) but did reveal weak homology between the N-terminal portion of region 2 (residues 109-167) and a region of the vertebrate proto-oncogene transcription factor c-MYC (Schmidt, 1999).

Furthermore, this region of c-myc lies in the linker between the DNA-binding domain and the transcriptional activation domain (Kerkhoff and Bister, 1991; Classon et al., 1993) and is not required for the oncogenic transformation activity of the protein (Stone et al., 1987). By analogy, this portion of region 2 of VRN1 may similarly serve as a linker region of no great importance to VRN1 function. Alternatively, region 2 may function as a novel type of transcriptional activation or repression domain, or in some other, unknown, function of VRN1. Table 4 gives information on the sequences which were used in comparisons with VRN1. The RTV1 gene is discussed below.

*(B) Effect of the allelic mutations on VRN1*

15

The mutations found in the two mutant alleles of *vrn1* (Figure 6) and the effect of these mutations on the resulting encoded protein can be correlated with the phenotypic severity, i.e., effect on vernalization response (Figure 1A), of the two alleles. As shown in Figure 7, the *vrn1-1* allele encodes a polypeptide of only 47 aa, and the *vrn1-2* allele encodes a polypeptide of 194 aa (the last six of which are incorrect due to a frameshift) compared to 341 aa for the wild-type protein. The fact that the polypeptide encoded by *vrn1-2* contains the first B3 domain as well as the putative PEST regions and NLS but is only slightly less severe in its effect on vernalization than the *vrn1-1* allele (Figure 1A), suggests that the second B3 DNA-binding domain may be required (but not necessarily sufficient) for VRN1 function under the conditions used.

30

*(C) The RTV1 gene, a relative of VRN1*

Despite the presence of many plant proteins that contain the B3 domain, only one putative protein sequence has been found

35



with a domain structure identical to VRN1, i.e., containing regions 1-3 in the same configuration and with no additional domains. The gene encoding this protein, which is represented in the *Arabidopsis* EST database (Table 4), has been named RTV1 (related to VRN1). RTV1, which is on IGF BAC clone F13F21 on chromosome 1, encodes a protein of 301 aa which is very similar to VRN1 (Table 5). While the overall similarity between RTV1 and VRN1 is 74% (within the coding region), the similarity is greatest at the C-terminal end, with region 3 of RTV1 and VRN1 being 99% similar (Table 5). Outside of the coding region (i.e., in the UTRs, promoter region and introns), the VRN1 and RTV1 genes appear to be unrelated. However, the intron/exon organization of the RTV1 gene is similar to VRN1 and therefore the two genes are likely to be the result of a duplication event. The most notable difference between VRN1 and RTV1 is the deletion of 33 amino acids in the first B3 domain of RTV1. It is worth noting that this deletion does not affect the C-terminal, most conserved, portion of the B3 domain.

20

The finding of a gene that is very closely related to VRN1 suggests that RTV1 may serve a function in vernalization response or other aspect of flowering time control. Since the *vrn1-1* allele encodes a short polypeptide of only 47 amino acids, with no complete putative DNA-binding domain (Figure 7) it is likely to encode a non-functional polypeptide. The fact that *vrn1-1* mutant plants still retain a small response to vernalization (Figure 1) suggests the presence of other *Arabidopsis* factors that can partially substitute for VRN1 function. Since RTV1 is a closely-related VRN1 paralogue in *Arabidopsis* it may be such a factor.

25  
30

Another factor that may possibly be responsible for this functional redundancy is VRN2 (Chandler et al., 1996). Like

*vrn1-1 fca-1* mutants, *vrn2-1 fca-1* mutants also retain a partial response to vernalization, but *vrn1-1 vrn2-1 fca-1* triple mutants do not (data not shown). If it is assumed that *vrn1-1* is a "null" mutation, then this result suggests that VRN1 and VRN2 act in separate and partially redundant vernalization-promoting pathways.

Example 9 - Detection and isolation of VRN1-related genes from other plant species

A high-stringency Southern blot of genomic DNA from various cereals, when hybridized with a probe corresponding to the VRN1 transcript, specifically detected VRN1-related genes in millet (FINGER, FOXTAIL 863B-PEARL, 841B-PEARL), sorghum (P20, P10), barley (BETZER, TRIUMPH, IGRI), rice (63-83, IR20), wheat (SYNTHETIC, SQL, CHINESE SPRING), and maize (P9, P10, C0, C8, C2, C9, DPT A, DTP79, B73, M017, B84, 12B84, L175, L25, DTP A, DW) (results not shown).

To prepare the blot, approximately 10 µg of genomic DNA from each of these varieties was digested with Eco RI (37°C, overnight). DNA samples were separated by gel electrophoresis on a 0.8% agarose gel run at 50V for 16 hours. The gel was then processed for Southern blotting by standard procedures (see Maniatis, supra) and DNA was blotted overnight onto a nylon membrane (Hybond-N, Amersham). Following blotting, the DNA was cross-linked to the filter by exposure to UV light according to the manufacturer's recommendations and baked at 80°C for 2 hrs.

The VRN1 cDNA probe V2V6 was prepared by amplifying an aliquot of the first-strand cDNA synthesis from total RNA of *Arabidopsis* seedlings with the oligonucleotide primers V2 and V6. The resulting PCR product was purified by agarose gel

electrophoresis and labeled with  $^{32}\text{P}$ -dCTP by the random hexamer priming method (see Maniatis, supra).

Hybridisation of the filter with the radiolabelled probe, and  
5 subsequent washes, were under standard high stringency  
conditions using buffer comprising 5 x SSC, 5 X Denhart's  
solution, and 0.5% SDS at 65°C for 16 hours. The filter was  
then washed sequentially in (1) 2XSSC, 0.1% SDS at room  
temperature for 10 minutes; (2) 1XSSC, 0.1% SDS at 65°C for 15  
10 minutes; and (3) 0.1XSSC, 0.1% SDS at 65°C for 10 minutes.

The washed filter was exposed to a PhosphorImager plate  
(Molecular Dynamics) for 3 days prior to visualisation.

15 In the light of the results above, in addition to the  
monocots, it is highly probable that *VRN1*-related genes will  
be found to exist in agronomically important dicot species  
(e.g. Brassicaceae, sugarbeet, peas and celery etc.)

20 Thus the provision of sequence information for the *VRN1* gene  
of *Arabidopsis thaliana* enables the obtention of homologous  
sequences from cDNA or genomic libraries from other plant  
species, such as can be prepared or obtained by the skilled  
person without undue burden. Positive clones can be further  
25 analyzed by restriction endonuclease digestion and Southern  
blotting as described hereinbefore. Particularly preferred  
are homologues from commercially important species that have a  
vernalization requirement, or show some response to  
vernalization.

30

#### Materials and Methods used in Examples

##### *Plant growth*

For vernalization treatments, seeds were sown on fine grit (Levington's M3) in individual pots, and germinated for increasing durations at 4°C, 8hr light:16hr dark, 5 mmol m<sup>-2</sup> sec<sup>-1</sup> light intensity. For dose-response experiments seed sowing was staggered, with all plants removed from the vernalization conditions simultaneously. Following vernalization, seedlings were placed into a controlled environment chamber (Gallenkamp), 20°C, 16 hr light: 8hr dark 90 mmol m<sup>-2</sup> sec<sup>-1</sup> light intensity. Seedlings receiving no vernalization treatment were stratified for 2 days under vernalization conditions, and grown for two days prior to transfer in to the growth cabinet. Plants were grown for 10 days, and then pricked out into individual compartments of P40 trays. Flowering time was measured by counting total leaf number (i.e. rosette and cauline leaves) by marking the leaves with permanent black ink as they emerged.

#### *Genetic mapping*

VRN1 was initially positioned on Chromosome 3 through linkage to RFLP markers mi339 and mi207 (Liu et al., 1996), in F2 progeny (154 chromosomes) of a cross between *vrn1-1 fca-1* (Ler background) and *fca-10* (Ws background), as described in (Chandler et al., 1996). As a first step in refining this map position, two existing RFLP markers in the region (g4711 and m560B2; Chang et al., 1988), and two existing SSLP markers in the region (ATHCHIB and ngal62; Bell and Ecker, 1994), were scored on a larger F2 population (988 chromosomes) of the same cross as above. In order to refine the map position of VRN1 further, new genetic markers that were polymorphic between Ler and Ws were developed (Table 1). Standard techniques (e.g., restriction digestion, <sup>32</sup>P-labeling of probes, agarose gel electrophoresis, Southern blotting, and PhosphorImager detection) were used throughout.

*Physical mapping*

YAC, BAC and cosmid clones and libraries were handled, analyzed, and hybridized according to standard procedures (Schmidt and Dean, 1995; Bent et al., 1998; Macknight et al., 1997). As with the genetic mapping of *VRN1*, some probes and PCR markers were existing and available, and some were developed in order to establish or refine the overlap between clones. The following probes and PCR markers were existing and available: mi289, GBGe303, MSH2, ve039, mi339, agp14, MAP2K, sAT2105b, and m506B2. New probes and PCR markers developed in order to identify the *VRN1* gene are listed in Table 2. New probes and PCR markers were developed by three methods: (1) iPCR of BAC ends, (2) design of PCR primers based on BAC end sequence data (from TIGR;website address:www.tigr.org/tdb/at/atgenome/bac\_end\_search/bac\_end\_search.html), and (3) sequencing of cosmid ends and design of PCR primers based on the obtained data.

*(A) iPCR of pBelo-BAC ends*

The following procedure is a modification of a protocol received from T. Altmann (MPI, Golm, Germany). DNA from 1/10th of a 25 ml BAC overnight culture was digested with (1) HhaI or EcoRI or HincII or RsaI for the T7 end, or (2) HhaI or HaeII or EcoRV for the Sp6 end, and phenol chloroform extracted and ethanol precipitated. Digested material was ligated in a 100 µl standard reaction with T4 DNA ligase, heat inactivated, and ethanol precipitated. Ligation products were digested with PvuI for the T7 end, and BsrBI for the Sp6 end in a 15 µl reaction volume. For PCR, 1 µl of digestion reaction was amplified in a standard reaction using (1) primers BACT7U and BACT7L for the T7 end, or (2) primers Sp6A and Sp6B for the Sp6 end.

BACT7U      5'- CCTCTTCGCTATTACGCCAG -3'  
BACT7L      5'- GCCCTTCCCAACAGTTCG -3'  
Sp6A        5'- CACACAGGAAACAGCTAT -3'  
Sp6B        5'- ACACAACATACGAGCCGGAA -3'

5

*(B) Sequencing of cosmid DNA and PCR products*

Genomic sequence was obtained from the ends of cosmid insert DNA using the BIGDYE cycle sequencing kit (Perkin Elmer Applied Biosystems), and T3 and T7 primers, whose sequences flank the genomic DNA insert site. For sequencing regions further into the cosmid insert DNA, and for sequencing PCR products amplified off of genomic DNA from the *vrn1-1* and *vrn1-2* alleles, the oligonucleotides shown in Table 3 were used. The reactions were run on an ABI377 machine, and compiled using the SeqMan (DNASTar, Lasergene) program.

*Cosmid complementation*

Cosmids in the 04541 binary vector were mobilized into *Agrobacterium tumefaciens* (strain C58C1 Rif<sup>R</sup>) by tri-parental mating (Hoekema et al., 1983). *vrn1-1 fca-1* plants were transformed with these *Agrobacterium* strains by root infection (Hooykaas, 1989). Transgenic plants were selected on GM with Kanamycin (50 mg/mL), and transferred to soil when they had reached the 3-4 leaf stage. The presence of each cosmid in the transgenic lines was confirmed using a specific diagnostic PCR reaction, using a primer present within the cosmid insert sequence and a primer present in the cosmid flanking the insert site. T2 seed were collected, and analyzed for the segregation of Kanamycin resistance or sensitivity on GM plates containing Kanamycin (as above), scored 14-20 days after germination. Lines that segregated a 3:1 ratio of resistant to sensitive plants were tested for their ability to

complement the *vrn1-1* mutant phenotype, by vernalizing for 5 weeks and recording the total leaf number.

#### *RT-PCR and 3'-RACE*

5

In order to determine the intron-exon structure of the *VRN1* gene, RT-PCR reactions using total RNA prepared from *fca-1* and *vrn1-1 fca-1* seedlings grown on soil were performed according to standard procedures (Frohman et al., 1988). The  
10 PCR products were sequenced using both the primers used for PCR, and selected internal primers, using the BIGDYE kit (PE Applied Biosystems). The reactions were run on an ABI377 machine, and compiled using the SeqMan (DNASTar, Lasergene) program.

15

#### *Sequence comparisons*

The nucleic acid sequence comparison in Table 5 was by using the Jotun Hein method (weighted residue table) of MegAlign  
20 (DNASTar). Genomic and cDNA sequences were aligned using the BLAST 2 SEQUENCES program (website address: [www.ncbi.nlm.nih.gov/gorf/bl2.html](http://www.ncbi.nlm.nih.gov/gorf/bl2.html)) from NCBI.

Parameters are preferably set, using the defaults, as follows:

25

Gap penalty: 11

Gap length penalty: 3

KTUP word length: 6

30 Amino acid sequences were initially aligned using the Clustal method, using the PAM 250 residue weight table, and further adjusted manually. For amino acid similarity comparisons, amino acids were grouped into five classes on the basis of physiochemical properties, as follows: (1) hydrophobic - G, A,  
35 V, P, M, I, L; (2) polar - S, T, N, Q, C; (3) bulky ring - Y,

F, W, H; (4) positively charged - K, R; (5) negatively charged  
- D, E.



Table 1. Genetic markers developed in order to identify the *VRN1* gene.

	Marker	Type	Ler/Ws polymorphism
5	GBGe303	RFLP	SspI; Ler band < Ws band
	MSH2	CAPS	Sau3A; Ler: 2 sites, Ws: no sites
			BfaI; Ler: no sites, Ws: 1 site
	ve039	CAPS	RsaI; Ler: 3 sites, Ws: 2 sites
	agp14	RFLP	HpaII; Ler band < Ws band
	pKS1240	RFLP	DraI; Ler band > Ws band
10	MAP2K	RFLP	HaeIII; Ler band < Ws band

Table 2. Physical mapping markers developed in order to identify the *VRN1* gene.

5	Marker	Type	Method used to develop
	T4L24-T7	Southern probe	iPCR
	T7H5-Sp6	Southern probe	iPCR
	T15C16-Sp6	Southern probe	iPCR
	T10N5-T7	Southern probe	iPCR
10	T24F13-T7	PCR diagnostic	BAC end sequence data (public)
	8H8-T7	PCR diagnostic	cosmid end sequence (obtained)
	F25C7-T7	PCR diagnostic	BAC end sequence data (public)
	10F10-T3	PCR diagnostic	cosmid end sequence (obtained)
	8H8-T3	PCR diagnostic	cosmid end sequence (obtained)
15	F18G1-T7	PCR diagnostic	BAC end sequence data (public)
	10F10-T7	PCR diagnostic	cosmid end sequence (obtained)
	T4L24-Sp6	PCR diagnostic	BAC end sequence data (public)
	T20A21-Sp6	Southern probe	iPCR
	T7H5-T7	Southern probe	iPCR
20	T15C16-T7	Southern probe	iPCR
	F10N5-Sp6	Southern probe	iPCR
	T24F13-Sp6	Southern probe	iPCR
	F28N8-Sp6	PCR diagnostic	BAC end sequence data (public)
	F5G10-T7	PCR diagnostic	BAC end sequence data (public)
25			

Table 3. Oligonucleotides developed to identify the *VRN1* gene. The positive (+) strand oligos correspond to the forward, or mRNA, strand of DNA, and the negative (-) strand oligos correspond to the reverse, or coding, strand of DNA.

5 The position indicated in the table refers to the nucleotide position in the *VRN1* genomic sequence (Annex I) of the 5' end of the oligo.

	Oligo	Strand	Position	Sequence (5' to 3')
10	S63	+	850	CAACGGTTAGCCCAAAC
	S64	-	866	GTTTGGGCTAACCGTTG
	V11	+	1193	GAGACCAGTTTGTTC
	S62	-	1229	GACAAATATAGGTGGAAAGG
	S66	+	1441	AAAGGGGAGTAGGTGGG
15	V7	+	1811	CTCTCTGGTCTTCTCTC
	V10	-	1828	GAAGAGAAGACCAGAGAG
	V6	+	1907	TTTTCTCATCCACTATCC
	S51	-	1930	TTTCTTGGATAGTGGATGAG
	S65	-	2166	AAAACAGGGAAGAGTAAGAAG
20	S52	+	2270	CATTGGTTGTGTTTGGTGGG
	V5	+	2599	GGTCTCTATGTATTGTGC
	V4	-	2616	GCACAATACATAGAGACC
	V12	-	2846	AGATTGATTACACGACTCC
	V8	+	3125	CCCAGATAAGTTTGTGAG
25	V3	+	3391	ATTCCGCTCACAACCAC
	V15	-	3414	GTTTGAAGTGGTTGTGAG
	V14	+	3477	TACCCATCACCCTTCC
	S60	-	3474	CAGAAGAAGGAAAGATGACC
	S61	+	3927	GAAGAAAGAGAGAGAGCC
30	V13	+	3976	ACCCTTTCTTCAGAGTG
	V9	-	3942	CTCTCTCTCTTCTTCTG
	V16	-	3993	CCACTCTGAAGAAAGGG
	S46	+	4096	CCTTCTGTTTCTGTTTCTC
	S45	-	4114	GAGAAACAGAAACAGAAGG
35	V2	-	4431	AAGATACTCCTACACGAC
	V17	+	4486	GTCTCGTTTTTCTCTCGG
	S49	-	4870	CTACCACAGTTCCCACCTAC

Table 4 Sequences corresponding to ESTs for *VRN1* and *RTV1*, and other sequences used for comparison to *VRN1*.

	Name	Type	Description	Accession #
5	92M2	nucl.	EST; <i>VRN1</i> transcript	T21005
	F2H7	nucl.	EST; <i>VRN1</i> transcript	N95889
	105022	nucl.	EST; <i>RTV1</i> transcript	T22671
	151H18	nucl.	EST; <i>RTV1</i> transcript	T76788
	247A13	nucl.	EST; <i>RTV1</i> transcript	AA713228
10	89H14	nucl.	EST; <i>RTV1</i> transcript	T20909
	89I23	nucl.	EST; <i>RTV1</i> transcript	T20917
	VRN1	aa	Encoded by <i>VRN1</i> (putative)	N/A
	RTV1	aa	Encoded by <i>RTV1</i> (putative)	N/A
15	3859591	aa	Putative <i>Arabidopsis</i> protein	AAC72857
	CAA19759	aa	Putative <i>Arabidopsis</i> protein	CAA19759
	CAA19755	aa	Putative <i>Arabidopsis</i> protein	CAA19755
	CAA19754	aa	Putative <i>Arabidopsis</i> protein	CAA19754
	RAV1	aa	Putative <i>Arabidopsis</i> protein	BAA34250
	FUSCA3	aa	Putative <i>Arabidopsis</i> protein	AAC35246
	ABI3	aa	Putative <i>Arabidopsis</i> protein	JQ1676
20	ARF1	aa	Putative <i>Arabidopsis</i> protein	AAD39318
	IAA24	aa	Putative <i>Arabidopsis</i> protein	AAB92476
	c-MYC	aa	Putative <i>Carassius</i> <i>auratus</i> (goldfish) protein	P49709

Table 5 Comparison of the Nucelotide and Amino Acid Sequences of *RTV1* to *VRN1*

5	Sequence	Nucleotide			Amino			
		Id	Length	Range <sup>b</sup>	Id	Sim.	Length	Range <sup>d</sup>
		a	(bp)		a	c	(aa)	
	RTV1	69	1026	269-	67	74	341	1-341
	complete			1291				
	RTV1 Region	49	283	272-550	42	44	93	2-94
	1							
10	RTV1 Region	71	429	551-981	71	82	144	95-238
	2							
	RTV1 Region	84	314	982-	95	99	103	239-341
	3			1291				

15 a Identity (%)

b Numbered relative to *VRN1* transcript sequence (Figure 7)

c Similarity (%)

d Numbered relative to *VRN1* encoded amino acid sequence (Figure 7)

20

## References

Abe, H., Yamaguchi-Shinozaki, K., Urao, T., Iwasaki, T., Hosokawa, D., and Shinozaki, K. (1997). Role of arabidopsis MYC and MYB homologs in drought- and abscisic acid- regulated gene expression. Plant Cell 9, 1859-68.

Abel, S., Oeller, P. W., and Theologis, A. (1994). Early auxin-induced genes encode short-lived nuclear proteins. Proc Natl Acad Sci U S A 91, 326-30.

Baker, S. S., Wilhelm, K. S., and Thomashow, M. F. (1994). The 5'-region of Arabidopsis thaliana cor15a has cis-acting

elements that confer cold-, drought- and ABA-regulated gene expression. *Plant Mol Biol* 24, 701-13.

Barnes, J. A., and Gomes, A. V. (1995). PEST sequences in calmodulin-binding proteins. *Mol Cell Biochem* 149-150, 17-27.

Becker, D., Brettschneider, R., and Lorz, H. (1994). Fertile transgenic wheat from microprojectile bombardment of scutellar tissue. *Plant J* 5, 299-307.

Bell, C. J., and Ecker, J. R. (1994). Assignment of 30 Microsatellite Loci to the Linkage Map of Arabidopsis. *Genomics* 19, 137-144.

Bent, E., Johnson, S., and Bancroft, I. (1998). BAC representation of two low-copy regions of the genome of *Arabidopsis thaliana*. *Plant J* 13, 849-55.

Camilleri, C., Lafleurriel, J., Macadre, C., Varoquaux, F., Parmentier, Y., Picard, G., Caboche, M., and Bouchez, D. (1998). A YAC contig map of *Arabidopsis thaliana* chromosome 3. *Plant J* 14, 633-42.

Chandler, J., Wilson, A., and Dean, C. (1996). *Arabidopsis* mutants showing an altered response to vernalization. *Plant Journal* 10, 637-644.

Chang, C., Bowman, J. L., DeJohn, A. W., Lander, E. S., and Meyerowitz, E. M. (1988). Restriction fragment length polymorphism linkage map for *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A* 85, 6856-60.

Chevallier, P. (1993). PEST sequences in nuclear proteins. *Int J Biochem* 25, 479-82.

Classon, M., Wennborg, A., Henriksson, M., and Klein, G. (1993). Analysis of c-Myc domains involved in stimulating SV40 replication. *Gene* 133, 153-61.

- 5 Cooper, K. F., Mallory, M. J., Smith, J. B., and Strich, R. (1997). Stress and developmental regulation of the yeast C-type cyclin Ume3p (Srb11p/Ssn8p). *Embo J* 16, 4665-75.

10 Frohman, M. A., Dush, M. K., and Martin, G. R. (1988). Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. *Proc Natl Acad Sci U S A* 85, 8998-9002.

15 Gilmour, S. J., and Thomashow, M. F. (1991). Cold-acclimation and cold-regulated gene-expression in ABA mutants of *Arabidopsis thaliana*. *Plant Molecular Biology* 17, 1233-1240.

20 Giraudat, J., Hauge, B. M., Valon, C., Smalle, J., Parcy, F., and Goodman, H. M. (1992). Isolation of the *Arabidopsis thaliana* ABI3 (Absciscic acid-insensitive) gene by positional cloning. *The Plant Cell* 4, 1251-1261.

25 Goldsbrough, A. P., Albrecht, H., and Stratford, R. (1993). Salicylic acid-inducible binding of a tobacco nuclear protein to a 10 bp sequence which is highly conserved amongst stress-inducible genes. *Plant J* 3, 563-71.

30 Gomes, A. V., and Barnes, J. A. (1997). Protein phosphatases are pest containing proteins. *Biochem Mol Biol Int* 41, 65-73.

Guilfoyle, T. J., Ulmasov, T., and Hagen, G. (1998). The ARF family of transcription factors and their role in plant hormone- responsive transcription. *Cell Mol Life Sci* 54, 619-27.

Hoekema, A., Hirsch, P. R., Hooykaas, P. J. J., and Schilperoort, R. A. (1983). A binary plant vector strategy based on separation of vir- and T-region of the *Agrobacterium tumefaciens* Ti-plasmid. *Nature* 303, 179-180.

5

Hooykaas, P. J. (1989). Transformation of plant cells via *Agrobacterium*. *Plant Mol Biol* 13, 327-36.

10

Jiang, C., Iu, B., and Singh, J. (1996). Requirement of a CCGAC cis-acting element for cold induction of the BN115 gene from winter *Brassica napus*. *Plant Mol Biol* 30, 679-84.

15

Kagaya, Y., Ohmiya, K., and Hattori, T. (1999). RAV1, a novel DNA-binding protein, binds to bipartite recognition sequence through two distinct DNA-binding domains uniquely found in higher plants. *Nucleic Acids Res* 27, 470-8.

20

Kerkhoff, E., and Bister, K. (1991). Myc protein structure: localization of DNA-binding and protein dimerization domains. *Oncogene* 6, 93-102.

25

Kim, J., Harter, K., and Theologis, A. (1997). Protein-protein interactions among the Aux/IAA proteins. *Proc Natl Acad Sci U S A* 94, 11786-91.

30

Koepp, D. M., Harper, J. W., and Elledge, S. J. (1999). How the cyclin became a cyclin: regulated proteolysis in the cell cycle. *Cell* 97, 431-4.

Laney, J. D., and Hochstrasser, M. (1999). Substrate targeting in the ubiquitin system. *Cell* 97, 427-30.

Liu, Y. G., Mitsukawa, N., Lister, C., Dean, C., and Whittier, R. F. (1996). Isolation and mapping of a new set of



129 RFLP markers in *Arabidopsis thaliana* using recombinant inbred lines. *Plant J* 10, 733-6.

Liu, Z. P., Galindo, R. L., and Wasserman, S. A. (1997). A  
5 role for CKII phosphorylation of the cactus PEST domain in dorsoventral patterning of the *Drosophila* embryo. *Genes Dev* 11, 3413-22.

Loake, G. J., Faktor, O., Lamb, C. J., and Dixon, R. A.  
10 (1992). Combination of H-box [CCTACC(N)7CT] and G-box (CACGTG) cis elements is necessary for feed-forward stimulation of a chalcone synthase promoter by the phenylpropanoid-pathway intermediate p-coumaric acid. *Proc Natl Acad Sci U S A* 89, 9230-4.

15 Luerksen, H., Kirik, V., Herrmann, P., and Misera, S. (1998). FUSCA3 encodes a protein with a conserved VP1/AB13-like B3 domain which is of functional importance for the regulation of seed maturation in *Arabidopsis thaliana*. *Plant J* 15, 755-  
20 64.

Macknight, R., Bancroft, I., Page, T., Lister, C., Schmidt, R., Love, K., Westphal, L., Murphy, G., Sherson, S., Cobbett, C., and Dean, C. (1997). *FCA*, a gene controlling flowering  
25 time in *Arabidopsis*, encodes a protein containing RNA-binding domains. *Cell* 89, 737-745.

Mannervik, M., Nibu, Y., Zhang, H., and Levine, M. (1999).  
30 Transcriptional coregulators in development. *Science* 284, 606-9.

Marchal, C., Haguenauer-Tsapis, R., and Urban-Grimal, D. (1998). A PEST-like sequence mediates phosphorylation and efficient ubiquitination of yeast uracil permease. *Mol Cell*  
35 *Biol* 18, 314-21.

Maule, J. (1997). Physical mapping by pulsed-field gel electrophoresis. *Methods Mol Biol* 68, 93-121.

McCarty, D. R., Hattori, T., Carson, C. B., Vasil, V., Lazar, M., and Vasil, I. K. (1991). The Viviparous-1 developmental gene of maize encodes a novel transcriptional activator. *Cell* 66, 895-905.

McKinsey, T. A., Chu, Z. L., and Ballard, D. W. (1997). Phosphorylation of the PEST domain of IkappaBbeta regulates the function of NF-kappaB/IkappaBbeta complexes. *J Biol Chem* 272, 22377-80.

Mhiri, C., Morel, J. B., Vernhettes, S., Casacuberta, J. M., Lucas, H., and Grandbastien, M. A. (1997). The promoter of the tobacco Tnt1 retrotransposon is induced by wounding and by abiotic stress. *Plant Mol Biol* 33, 257-66.

Mozo, T., Fischer, S., Shizuya, H., and Altmann, T. (1998). Construction and characterization of the IGF Arabidopsis BAC library. *Mol Gen Genet* 258, 562-70.

Nordin, K., Vahala, T., and Palva, E. T. (1993). Differential expression of two related, low-temperature-induced genes in *Arabidopsis thaliana* (L.) Heynh. *Plant Mol Biol* 21, 641-53.

Rechsteiner, M., and Rogers, S. W. (1996). PEST sequences and regulation by proteolysis. *Trends Biochem Sci* 21, 267-71.

Rogers, S., Wells, R., and Rechsteiner, M. (1986). Amino acid sequences common to rapidly degraded proteins: the PEST hypothesis. *Science* 234, 364-8.

Schmidt, E. V. (1999). The role of c-myc in cellular growth control. *Oncogene* 18, 2988-96.

Schmidt, R., and Dean, C. (1995). Hybridization analysis of YAC clones. *Methods in Mol. & Cell. Biol.* 5, 309-318.

Steger, D. J., Utley, R. T., Grant, P. A., John, S.,  
5 Eberharther, A., Cote, J., Owen-Hughes, T., Ikeda, K., and  
Workman, J. L. (1998). Regulation of transcription by  
multisubunit complexes that alter nucleosome structure. *Cold  
Spring Harb Symp Quant Biol* 63, 483-91.

10 Stockinger, E. J., Gilmour, S. J., and Thomashow, M. F.  
(1997). *Arabidopsis thaliana* CBF1 encodes an AP2 domain-  
containing transcriptional activator that binds to the C-  
repeat/DRE, a cis-acting DNA regulatory element that  
stimulates transcription in response to low temperature and  
15 water deficit. *Proc Natl Acad Sci U S A* 94, 1035-40.

Stone, J., de Lange, T., Ramsay, G., Jakobovits, E., Bishop,  
J. M., Varmus, H., and Lee, W. (1987). Definition of regions  
in human c-myc that are involved in transformation and  
20 nuclear localization. *Mol Cell Biol* 7, 1697-709.

Suzuki, M., Kao, C. Y., and McCarty, D. R. (1997). The  
conserved B3 domain of VIVIPAROUS1 has a cooperative DNA  
binding activity. *Plant Cell* 9, 799-807.

25 Thomashow, M. F. (1994). *Arabidopsis thaliana* as a model for  
studying mechanisms of plant cold tolerance. In *Arabidopsis*  
(New York: Cold Spring Harbor Laboratory Press), pp. 807-834.

30 Torchia, J., Glass, C., and Rosenfeld, M. G. (1998). Co-  
activators and co-repressors in the integration of  
transcriptional responses. *Curr Opin Cell Biol* 10, 373-83.

Ulmasov, T., Hagen, G., and Guilfoyle, T. J. (1997). ARF1, a transcription factor that binds to auxin response elements. *Science* 276, 1865-8.

- 5 Urao, T., Yamaguchi-Shinozaki, K., Urao, S., and Shinozaki, K. (1993). An Arabidopsis myb homolog is induced by dehydration stress and its gene product binds to the conserved MYB recognition sequence. *Plant Cell* 5, 1529-39.
- 10 Vierstra, R. D. (1996). Proteolysis in plants: mechanisms and functions. *Plant Mol Biol* 32, 275-302.
- Yaglom, J. A., Goldberg, A. L., Finley, D., and Sherman, M. Y. (1996). The molecular chaperone Ydj1 is required for the
- 15 p34CDC28-dependent phosphorylation of the cyclin Cln3 that signals its degradation. *Mol Cell Biol* 16, 3679-84.

Annex I - Ler VRN1 genomic (contig 29 [1501-6500])

	1	10	20	30	40
	TTTAAAATTCGAATTGGGATTTAAGAAAAATTCTCATCAA				
5	ATATTTATCATTAGTGTATATATATCAGTGTTTACATTT				
	GTTAATCCTAAATAATAAACCGATCTGAAAAGTTGATAAA				
	TGCGTTGTCAAAGACAAAATATACATCCAAACAAATCAC				
	GTGATTGCCTTCAACTTGCCACgGGTTCAAAGATTTAACA				
	AATCTTCTAAAACACCAACTTAACCCACGAATACACAAGC				
10	ACAGAGTGGTGGTAAACATACAAGTTAATGAGTTATTCAA				
	ATGAGATTTTCAATATCATTCCTTCTTCAGCCCGTCACAAG				
	AAGCCAAGATTAAGCCATTAGAGGAAGTTTATAAACCGAC				
	AAAACCTGCTTAGATACAAAGAATACTAGCTAATGTGTTT				
	CAACAACTTCAAATTGACGATACGTTACATTCATATTAA				
15	TCACTTCAGAGCTTGATTATTCAAATTATTTTTTCTACTG				
	TGATACATATATACACACATGTTTTGCTTTTCTATGATTC				
	TATCTACATTTTCATACCGTTGAATAATTTATGTATGAAT				
	TACGATGCAATTTTCCTTCATTATGCTTGAATAAAATGCTT				
	TTGGACATGCATGCGATATTGGATCTACTTTTGGATTCTA				
20	TTTTTTAAAATCAGCGAGTTTGTTGCTTTGTAATTTTTTAA				
	TTAGGCATCAAGAATTTCTAAAATGCACGCGAACTGGTGA				
	AAAGAGGAATGTTTACGTTTACCCCTTTATTTTCTTACAG				
	CTCATAAGGATACTGTCAGAAGACAGAACCAAGGCTCTCT				
	GACTATAAATTTGGAATCCATTTAAACATAATGTTATGACC				
25	AATGATGGCCAACGGTTAGCCCAAATAATTAACATAAAG				
	TCAAGTTCCAATATTCTAAGGAGAAATAATAGTATACTAA				
	ACATACATTAGAGAGGTTAAACTTCTTTTTGGATTTAAGT				
	GTGTATGCATAGGCTATTTATTCTTAAGTATAACTATTAA				
	CTGTAGCTAGATTTATACAAGAAATACATAAAACTTTATG				
30	CATGTGAGGTAGCCATGAATATACGTACATGTTGCAATCG				
	ATTATACATGTTGTATTTGGATTTCTCTATACATGTTTTA				
	ACTTGTCATTCTCTAAGTATATACATACCATTAATACTGT				
	GGGCATGAGTTTATGATAAGACTTTTCTTTTGGAGACCAG				
	TTTTGTTTTCTTTCCACCTATATTTGTCTATAGGCTTCa				
35	gACGGTACACTAGTTTACAAGTGTTTTTATATGTTCTAAA				

TAAAATTGAGATTTTCCGGAACGGTATGATCTGTTTGCAA  
ATAAGGACGTATATATAACAGTATCAAATATATTTGTTGT  
TATAAGGCAATAATATATTTTCTGAGATATTGCGTGTTAC  
AAAAAAGAAATATTTGTTAAGAAAAAAAAAAGATGGTCGAA  
5 AAAGGGGAGTAGGTGGGGGCGGTCGGCTTTTGATTAGTTA  
ATAAAAGAAACCACACGAGTGACCTACCGATTGCACTCAA  
CGAGTCTACCGAGCTAACACAGATTCAACTCGCTCGAGCT  
TCGTTTTATGACAAGTTGGTTTTTTTTTTTTTTTTTAAT  
TTTTTCATCTTCTTGGGTTTGGTTGGGTCACTCTTCAGGT  
10 CAGGTGTGTAAAAAAGAAAGAAAGAAAGAGAGATTGTTG  
TGTTGTAACCCCTTTGACTAAAATCTAATGAACTTTTTTA  
ACACAACAAAACCTCCTTCAGATCTGAAAGGGTCTTCTTC  
TCTCTTAGTCTCTTTGTCCTTTTATTCTCCGTCGTCGTTT  
CATGATCTGACTCTCTGGTCTTCTCTTCTTCTTCTTCTTC  
15 TTCTATTTTTTCTTACTTCGTCACTGTTGTGTCTGAACAT  
GCCACGCCCTTTCTTCCATAAGTTGATTTTCTCATCCACT  
ATCCAAGAAAAACGTCTGGTAACTTACTCTCTCTCTCTCT  
CTCTCTCTCTGTTCTCCTTCTCCTCATCTTTCAAAGTTTT  
GATTTTGTGCGAAATTGAGGGTTTTCAAGGTTTGGAATCT  
20 GGTGAACGAGTTTGTAAGATTATGCCTTGTGACACTCTTG  
CTTGATTTCTTACAATTCACTTGTATTGATTCTTTGTAAG  
AATCGAGTCAAGGTTGTGCTTTTATCTTCTTACTCTTCCC  
TGTTTTGGGTAATGAAAAGAAGTTCCATTTTTGAACTTTG  
TGTTGTCTTATTGGTCAAATGAGAATTGTGGGTTTCCAA  
25 TGGAAGTCTGCAAGACAGTTTCTTTTGGTCATTGGTTGTG  
TTTGGTGGGAAATTGGGTATTTGATGGTATATCTGTACTC  
TGACAGCATATTGTGTGTAGTTTGGGAATTTTTTTTTTTT  
TTTTGAGTGATTTGACTTTTGGAGGACGATTTGATTCTGT  
CAGATTGATCAAATTTCTTCTGAGGAGAAAAAGTTGAGAT  
30 CTGTTTATGGTTTCTCTATTATAAATGTCTGTTTTGTTTA  
CTCTATTTTGACTGTTTTCTCTGTTTGACTTAGGAATGTC  
TGAGATCTTAGACTCCTTATTGAGTATTGTGtGGCTTGTG  
AGTGAATCCCTAAACTGAGTAGTTGACTTGTTTTGAAGG  
TCTCTATGTATTGTGCTTATGTTTTAAAGTTGTCTACTTT  
35 ATTTGATACAGTGATTAGTCATCACTTGTACAGATTCCCC

CAAGAGCATTGTTTTGAACAAATCCAAATTTGCTTAGCTC  
TCCATTTGGCATTTAAGTGA TAGATTTTCTCTGGAATAA  
TGATTTTCGATTAACACAGGCATTTATGTGGAACCAAGTTT  
GCAAATTATTAATGTGATAAGATCATAGGAGTCGTGTAAT  
5 CAATCTATTCAGAGATAAATGTACCATTTTACATGTGTAC  
TAATGGACTGTGTCTCCTTGTTGATGCCTTCTCTAAACTG  
AAATATGGCCTTTTGGTTTGTGTTTTTAAATTAGGTAAAG  
CCGTCGTTTCTTCAGCTACTGTGTTTATTGGATGTTTTTG  
CTGAAAAATGTCTGTTTCGATTTGATGTTCTCGCAATATT  
10 CTGTGCTGTTCTTATAGATATTGTGGACATTTATATCATT  
ATATGCTTCTTTATATCTCATACCGGCATGCTTGTGCAGA  
GGGTCCCAGATAAGTTTGTGAGTAAATTCAAGGATGAGCT  
TTCGGTTGCTGTTGCACTCACAGTACCTGATGGTCATGTT  
TGRCGTG TAGGACTAAGGAAAGCTGACAACAAAATTTGGT  
15 TTCAAGATGGTTGGCAAGAGTTTGTTGACCGTTACTCCAT  
TCGCATTGGTTATCTTTTGATTTTTAGATATGAAGGAAAC  
TCTGCCTTCAGCGTCTACATTTTCAATTTATCCCACTCTG  
AGATCAATTACCATTCCACCGGTCTCATGGATTCCGCTCA  
CAACCACTTCAAACGCGCCCGTTTGTTTGAAGACCTTGAA  
20 GATGAAGATGCCGAGGTCATCTTTCCTTCTTCTGTGTACC  
CATCACCACCTCCTGAGTCTACAGTACCAGCCAACAAAGG  
GTATGCTAGTTTCAGCCATCCAAACCTTGTTCACTGGACCA  
GTTAAAGGTGATATTTATAACCAACTGATTCCCTTTATCT  
ATCGCTGATTACGCGTCTTATCATTCTTTTGAGGTTGATG  
25 CTTGATATTTTCCTTATCTCCAGCTGAAGAGCCAACGCCA  
ACCCCAAAAATACCTAAAAAGAGAGGGAGGAAGAAGaAAA  
ATGCTGATCCTGGTAAGCACTTTTCCTCTTTGAAATGCTT  
CAGACTCGTTTTTCAGAGGATTCACAGATTCTTCCTCATGA  
TACATATATCCTTTTGATATTGTCCTTACAGAGGAAATAA  
30 ACTCATCAGCTCCGCGAGATGATGATCCAGAGAACCGTTC  
AAAGTTCTACGAGAGTGCTTCTGCGAGAAAGAGAACCGTG  
ACTGCAGAAGAAAGAGAGAGAGCCATCAATGCAGCCAAAA  
CGTTCGAACCAACAAACCCCTTTCTTCAGAGTGGTTCTGCG  
ACCATCCTATCTATACAGAGGTTGCATCATGGTAATAAAA  
35 AAACATCTTAGGAAGACTTAATCTTATCGGTGTCTTCACT

GATCTCTAAAAGAAGCCTTCTGTTTCTGTTTCTCTCAACA  
GTATCTTCCTTCTGGGTTTGCTGAGAAGTACCTAAGTGGG  
ATCTCCGGGTTTCATCAAAGTCCAGCTTGCGGAGAAACAAT  
GGCCTGTTCGATGTCTCTACAAAGCCGGGAGAGCCAAATT  
5 CAGTCAAGGATGGTACGAATTCAGTCTAGAGAACAACCTTA  
GGAGAAGGAGACGTCTGTGTGTTTGAGCTGCTCAGAACCA  
GAGATTTTCGTTTTTGAAAGTGACAGCCTTTCGAGTCAACGA  
GTACGTCTGAACAAAGCATTATGGTGTGATCATTCTGGAT  
TTGCAAGTACAATGTCGTGTAGGAGTATCTTAATTTAAAA  
10 ACAACTAAAAAACTCTCTTCTGGTCTGTGTCATTATTGCG  
TCAGTGTCTCGTTTTTCTCTCGGGTTTACTTTGTGTTAT  
CGATGTGGATAAGTTGTTTTTACCTCATTATATAAACCT  
CTTGAGTGGAACTCAAATTGTTTGAGTAGAACAAACAAAG  
TTAGGGTTTAAGAAGAAGTCTGTAAATACCTAATCTCCAT  
15 CAAATTTGAGTAGAAAGACAAACTGTTCTGGTGGGAATACA  
AGGAGGGAACTTGAGATAACAACTTAAGAATAGCCTTCA  
AGCCAACGTCTAGAATTTGATGAAGTTGTTGTTTGATCAC  
CTCTGAGATAATTGGAAACCCTCTTCATGCAGTTTGCTTG  
AGGATACTGGTGAAAATGGGAGTATTGAAGGAAAATGCAT  
20 ATATAAGATTGTAGGTGGGAACTGTGGTAGCAGACACAAC  
ACTTGTTCTCTAGACATATACTGTACCAGACATGTTTTGA  
TCATAAAACTTAAAAAAAAGAAAACCGTGTGTAAATCAAG  
CAAGGAACAACTACAATATTACAATCTTATTGAGATATCA



CLAIMS

- 1 An isolated nucleic acid molecule which comprises a VRN1 nucleotide sequence encoding a polypeptide which is capable of specifically altering the vernalisation response of a plant into which the nucleic acid is introduced and expressed.
- 2 A nucleic acid as claimed in claim 1 wherein the VRN1 nucleotide sequence:
  - (i) encodes the VRN1 polypeptide of Fig 7, or
  - (ii) encodes a variant resistance polypeptide which is a homologous variant of the resistance polypeptide shown in Fig 7 and which shares at least about 50%, 60%, 70%, 80% or 90% identity therewith,
- 3 A nucleic acid as claimed in claim 1 or claim 2 wherein the VRN1 nucleotide sequence is that shown in Fig 7 from nucleotides 269-1295 inclusive, or a sequence which is degeneratively equivalent thereto.
- 4 A nucleic acid as claimed in claim 1 or claim 2 wherein the VRN1 nucleotide sequence is shown in Annex I.
- 5 A nucleic acid as claimed in claim 1 or claim 2 wherein the VRN1 nucleotide sequence encodes a derivative of the resistance polypeptide shown in Fig 7 by way of addition, insertion, deletion or substitution of one or more amino acids.
- 6 A nucleic acid as claimed in claim 1 or claim 2 wherein the VRN1 nucleotide sequence consists of an allelic or other homologous variant of the nucleotide sequence of claim 3

7 A nucleic acid as claimed in claim 6 wherein the VRN1 nucleotide sequence is the VRN1 paralogue RTV1 of Figure 9.

8 An isolated nucleic acid which comprises a nucleotide sequence which is the complement of the VRN1 nucleotide sequence of any one of the preceding claims.

9 An isolated nucleic acid for use as a probe or primer, said nucleic acid having a distinctive sequence of at least about 16-24 nucleotides in length, which sequence is present in Annex I or a sequence which is degeneratively equivalent thereto, or the complement of either.

10 A nucleic acid as claimed in claim 9 which is selected from the oligonucleotides (shown below in the 5' to 3' orientation):

S63	CAACGGTTAGCCCAAAC
S64	GTTTGGGCTAACCGTTG
V11	GAGACCAGTTTTGTTTTCC
S62	GACAAATATAGGTGGAAAGG
S66	AAAGGGGAGTAGGTGGG
V7	CTCTCTGGTCTTCTCTTC
V10	GAAGAGAAGACCAGAGAG
V6	TTTTCTCATCCACTATCC
S51	TTTCTTGGATAGTGGATGAG
S65	AAAACAGGGAAGAGTAAGAAG
S52	CATTGGTTGTGTTTGGTGGG
V5	GGTCTCTATGTATTGTGC
V4	GCACAATACATAGAGACC
V12	AGATTGATTACACGACTCC
V8	CCCAGATAAGTTTGTGAG
V3	ATTCCGCTCACAACCAC
V15	GTTTGAAGTGGTTGTGAG
V14	TACCCATCACCCTTCC
S60	CAGAAGAAGGAAAGATGACC
S61	GAAGAAAGAGAGAGAGCC
V13	ACCCTTTCTTCAGAGTG

V9 CTCTCTCTCTTTCTTCTG  
V16 CCACTCTGAAGAAAGGG  
S46 CCTTCTGTTTCTGTTTCTC  
S45 GAGAAACAGAAACAGAAGG  
V2 AAGATACTCCTACACGAC  
V17 GTCTCGTTTTTTCTCTCGG  
S49 CTACCACAGTTCCACCTAC  
8H8DIAG1 ACCTGCTTCTGCCAACCGCTC)

11 A process for producing a nucleic acid as claimed in claim 5 comprising the step of modifying a nucleic acid as claimed in claim 3 or claim 4.

12 A method for identifying or cloning a nucleic acid as claimed in claim 6 or claim 7, which method employs a nucleic acid as claimed in claim 9 or claim 10.

13 A method as claimed in claim 12, which method comprises the steps of:

- (a) providing a preparation of nucleic acid from a plant cell;
- (b) providing a nucleic acid molecule which is a nucleic acid as claimed in claim 9 or claim 10,
- (c) contacting nucleic acid in said preparation with said nucleic acid molecule under conditions for hybridisation, and,
- (d) identifying nucleic acid in said preparation which hybridises with said nucleic acid molecule.

14 A method as claimed in claim 12, which method comprises the steps of:

- (a) providing a preparation of nucleic acid from a plant cell;
- (b) providing a pair of nucleic acid molecule primers suitable for PCR, at least one of said primers being a nucleic acid as claimed in claim 9 or claim 10,

(c) contacting nucleic acid in said preparation with said primers under conditions for performance of PCR,  
(d) performing PCR and determining the presence or absence of an amplified PCR product.

15 A method as claimed in claim 14 wherein the preparation of nucleic acid is obtained from a Brassicaceae plant.

16 A recombinant vector which comprises the nucleic acid of any one of claims 1 to 8.

17 A vector as claimed in claim 16 wherein the nucleic acid is operably linked to a promoter for transcription in a host cell, wherein the promoter is optionally an inducible promoter.

18 A vector as claimed in claim 16 or claim 17 which is a plant vector.

19 A method for transforming a host cell, which comprises the step of introducing the vector of any one of claims 16 to 18 into a host cell, and optionally causing or allowing recombination between the vector and the host cell genome such as to transform the host cell.

20 A host cell containing or transformed with a heterologous vector of any one of claims 16 to 18.

21 A method for producing a transgenic plant, which method comprises the steps of:

- (a) performing a method as claimed in claim 19 wherein the host cell is a plant cell,
- (b) regenerating a plant from the transformed plant cell.

22 A transgenic plant which is obtainable by the method of claim 21, or which is a clone, or selfed or hybrid progeny or other descendant of said transgenic plant, which in each case includes a heterologous nucleic acid of any one of claims 1 to 8.

23 A plant as claimed in claim 22 which is selected from the list consisting of: rice; maize; wheat; barley; oats; rye; oil seed rape; sugar beet; maize; sunflower; soybean; sorghum; lettuce; endive; cabbage; broccoli; cauliflower; carnations; geraniums.

24 A part of propagule from a plant as claimed in claim 22 or claim 23, which in either case includes a heterologous nucleic acid of any one of claims 1 to 8.

25 An isolated polypeptide which is encoded by the VRN1 nucleotide sequence of any one of claims 1 to 7.

26 A polypeptide as claimed in claim 25 which is the VRN1 resistance polypeptide shown in Fig 4.

27 A method of making the polypeptide of claim 25 or claim 26, which method comprises the step of causing or allowing expression from a nucleic acid of any one of claims 1 to 7 in a suitable host cell.

28 A polypeptide which comprises the antigen-binding site of an antibody having specific binding affinity for the polypeptide of claim 26.

29 A method for assessing the vernalisation phenotype of a plant, the method comprising the step of determining the presence and/or identity of a VRN1 allele therein

comprising the use of a nucleic acid as claimed in claim 9 or claim 10.

30 A method for influencing or affecting the vernalisation phenotype of a plant, which method comprises the step of causing or allowing expression of a heterologous nucleic acid as claimed in any one of claims 1 to 8 within the cells of the plant, following an earlier step of introducing the nucleic acid into a cell of the plant or an ancestor thereof.

31 A method as claimed in claim 30 for modifying the kinetics and/or optimal temperature of the vernalization response such as to alter the phenotype of the plant with respect to any one or more of: geographic range; length of a vernalization period; length of a vegetative growth phase.

32 A method as claimed in claim 30 or claim 31 for reducing the vernalisation requirement of a plant, wherein the heterologous nucleic acid is that claimed in any one of claims 1 to 7.

33 A method as claimed in claim 30 or claim 31 for increasing the vernalisation requirement of a plant, which method comprises any of the following steps of:

- (i) causing or allowing transcription from a nucleic acid as claimed in claim 8 in the plant such as to reduce *VRN1* expression by an antisense mechanism;
- (ii) causing or allowing transcription from a nucleic acid as claimed in any one of claims 1 to 7 or a part thereof such as to reduce *VRN1* expression by co-suppression;
- (iii) use of nucleic acid encoding a ribozyme specific for a nucleic acid as claimed in any one of claims 1 to 7.

34 An isolated nucleic acid molecule encoding the promoter of the VRN1 gene, or a homologous variant thereof which has promoter activity.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
29 March 2001 (29.03.2001)

PCT

(10) International Publication Number  
WO 01/21822 A1

- (51) International Patent Classification<sup>7</sup>: C12N 15/82, C12Q 1/68, C07K 14/415, 16/16, A01H 5/00
- (74) Agents: KREMER, Simon, M. et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB).
- (21) International Application Number: PCT/GB00/03525
- (22) International Filing Date:  
13 September 2000 (13.09.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
9922071.7 17 September 1999 (17.09.1999) GB
- (71) Applicant (for all designated States except US): PLANT BIOSCIENCE LIMITED [GB/GB]; Norwich Research Park, Colney Lane, Norwich, Norfolk NR4 7UH (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): DEAN, Caroline [GB/GB]; 19 Waverley Road, Norwich, Norfolk NR4 6SG (GB). LEVY, Yaron, Yakov [US/DK]; RISO National Laboratory, Plant Biology & Biogeochemistry Department, Frederiksborgvej 399, Building 330, P.O. Box No.49, DK-4000 Roskilde (DK).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- With international search report.
  - Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/21822 A1

(54) Title: METHODS AND MEANS FOR MODIFICATION OF PLANT FLOWERING CHARACTERISTICS

(57) Abstract: Provided are isolated nucleic acid molecules which comprises VRN1 nucleotide sequences, which encode a polypeptide which is capable of specifically altering the vernalisation response of a plant into which the nucleic acid is introduced and expressed. Examples include cDNA and gDNA sequences (see e.g. Annex I). Also provided are variant molecules which may be derivatives or homologues (e.g. alleles, or paralogues such as *RTVI*), plus also complementary molecules. Corresponding polypeptides form a further part of the invention. The invention also provides methods and materials for preparing and using these molecules e.g. in the production of plants having modified vernalisation characteristics. Also methods for influencing and assessing the vernalisation phenotype of a plant.



Figure 1





Figure 2:




Figure 3





Figure 4:



10/088187

Figure 5





Figure 6



### Figure 7

	TCTTGGGTTTGGGTTGGGCACTCTTCAGGTTCAGGTGTTGAAAA	43
AA	GAAAGAAAGAAAAGAGAGATGTTGTGTGTGTAACCCCTTTGACTAAAAATCTAATGAACCTTTTAAACACAACA	118
AA	ACTCCTTCAGATCTGAAAGGGTCTCTCTCTCTCTTAGTCTCTTTGTCTCTTTATTCTCCGTCGTCGTTTCAT	193
GAT	CTGACTCTCTGGTCTTCTCTCTCTCTCTCTCTCTTCTTCTTCTTCTTACTTCGTCACTGTGTGTCTGAAC	268
AT	GCCACGCCCTTTCTTCCATAAGTTGATTTTCTCATCCACTATCCAAGAAAAACGTCTGAGGGTCCCAGATAAG	343
M	<u>P R P F F H K L I F S S T I Q E K R L R V P D K</u>	25
TT	TGTGAGTAAATTCAAGGATGAGCTTTTCGGTTGCTGTTGCACTCACAGTACCTGATGGTCATGTTTGCCGTGTA	418
F	<u>V S K F K D E L S V A V A L T V P D G H V (W) R V</u>	50
GG	ACTAAGGAAAGCTGACAAACAAATTTGGTTTCAAGATGGTTGGCAAGAGTTTGTGTACCGTTACTCCATTTCGC	493
G	<u>L R K A D N K I W F Q D G W Q E F V D R Y S I R</u>	75
ATT	GGTTATCTTTTGATTTTTAGATATGAAGGAAACTCTGCCTTCAGCGTCTACATTTTCAATTTATCCACTCT	568
I	<u>G Y L L I F R Y E G N S A F S V Y I F N L S H S</u>	100
GAG	ATCAATTACCATCCACCGGTCTCATGGATTCGCTCACAACCACCTTCAAACGCGCCCGTTTGTGTTGAAGAC	643
E	<u>I N Y H S T G L M D S A H N H F K R A R L F E D</u>	125
CT	TGAAGATGAAGATGCCGAGGTCTCTTTCTCTCTCTGTGTACCCATCACCACCTTCCTGAGTCTACAGTACCA	718
L	<u>E D E D A E V I F P S S V Y P S P L P E S T V P</u>	150
GC	CAACAAAGGGTATGCTAGTTTCAGCCATCCAAACCTTGTTCTACTGGACCAAGTTAAAGCTGAAGAGCCAACGCCA	793
A	<u>N K G Y A S S A I Q T L F T G P V K A E E P T P</u>	175
A	CCCCAAAATACCTAAAAAGAGAGGGAGGAAGAAAGAAAATGCTGATCCTGAGGAAATAAACTCATCAGCTCCG	868
T	<u>P K I P K K R G R K K K (N) A D P E E I N S S A P</u>	200
CG	AGATGATGATCCAGAGAACCCTTCAAAGTTCTACGAGAGTGCTTCTGCGAGAAAGAGAACCGTGACTGCAGAA	943
R	<u>D D D P E N R S K F Y E S A S A R K R T V T A E</u>	225
GAA	GAGAGAGAGCCATCAATGCAGCCAAAACGTTGCAACCAACAAACCCCTTTCTTCAGAGTGGTTCTGCGACCA	1018
E	<u>R E R A I N A A K T F E P T N P F F R V V L R P</u>	250
TC	TATCTATACAGAGGTTGCATCATGTATCTTCTCTCTGCGGTTTGCTGAGAAGTACCTAAGTGGGATCTCCGGG	1093
S	<u>Y L Y R G C I M Y L P S G F A E K Y L S G I S G</u>	275
TT	CATCAAAGTCCAGCTTTCGAGAGAAACATGGCCTGTTGATGTCTCTACAAAGCCGGGAGAGCCAAATTAGT	1168
F	<u>I K V Q L A E K Q W P V R C L Y K A G R A K F S</u>	300
CA	AGGATGGTACGAATTCACCTCTAGAGAACAACCTTAGGAGAAGGAGACGCTCTGTGTGTTTGTAGCTGCTCAGAAC	1243
Q	<u>G W Y E F T L E N N L G E G D V C V F E L L R T</u>	325
AG	AGATTTTCGTTTGAAGTGACAGCCTTTCGAGTCAACGAGTACGCTGGAACAAAGCATTATGGTGTGATCATT	1318
R	<u>D F V L K V T A F R V N E Y V</u>	341
CT	GGATTGCAAGTACAATGTCGTGTAGGAGTATCTTAATTTAAAAACAACATAAAAACTCTCTTCTGGTCTGTG	1393
TC	ATTATTGCGTCAGTGTCTCGTTTCTCTCTCGGGTTTACTTTGTGTTATCGATGTGGATAAGTTGTTTTACC	1468
TC	ATTATATGTAACCTCTTGAGTGGAA	1493

NLS (boxed), B3 DNA-binding domains (underlined), PEST regions (doubly underlined), protein kinase C phosphorylation site (asterisks). The positions of introns are indicated with arrowheads. The positions of the mutations in *vm1-1* and *vm1-2* are circled.

Figure 8

VRN1	MPRPFFHK---	20	LIFFSTIQEKRLRVDPKFKDEL	40	SVAVALTVPDGHVVRV	60	GLRKADN	57
RTV1	MPRSFFHMFNSL	20	LFLSSTQAS	40			GLRKANN	27
VRN1	KIWFQDGWQEFVDRYSIRIGYLLI	80	FRYEENSASFVYIEN--	100	ESHSEINYNHST--	120	GLM--D	111
RTV1	KIWFQDGWQEFVNRESIRIG	80	FRYKVM---	100	IVYIFQFYEPHSEINHSSSEALMQMD	120		79
VRN1	SAHNHF-KRARLFEDDE	140	DEDACVIEPSSVYPSPL	160	PESTVPANKGYA-SSA	180	IQRLFTGPVK	169
RTV1	SAONQENKRARLFEDDE	140	ELKDAKVIIYPSN-----	160	PESTEPEVANKGYGGS	180	TAIQSFKES-K	132
VRN1	AEEPTPTPKLKKRGRKKKNAD	200	PEEINSSAPRDD	220	PEENRSKFYESASARKRTVTAEERER	240		229
RTV1	ABE---TPKVLKKRGRKKKNEN	200	PEEVNSSTPGGDD	220	SENRSKFYESASARKRTVTAEERER	240		189
VRN1	AINAAKTFEPTNPFRVVL	260	RPSYLYRGCMYLP	280	SGFAEKYLSGISGFIK	300	VOLAEKOWPVR	289
RTV1	AVNAAKTFEPTNPFRVVL	260	RPSYLYRGCMYLP	280	SGFAEKYLSGISGFIK	300	VLGEKOWPVR	249
VRN1	CLYKAGRAKFSQGWYEFTLENN	320	EGGDV	340	CVFELLRTRDFVL		VTAFRVNEYV	341
RTV1	CLYKAGRAKFSQGWYEFTLENN	320	EGGDV	340	CVFELLRTRDFVL		VTAFRVNEYV	301



**DECLARATION, POWER OF ATTORNEY AND POWER TO INSPECT**

As below named inventor, I hereby declare:

that my residence, post office address and citizenship are as stated below next to my name;

that I verily believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the invention entitled: **METHODS AND MEANS FOR MODIFICATION OF PLANT FLOWERING CHARACTERISTICS** the specification of which [check one(s) applicable]

☒ was filed 13 September 2000 as International Patent Application No. PCT/GB00/03525, on which U.S. National Stage Application No. 10/088,187 is based; and/or  
☐ was amended by Amendment filed \_\_\_\_\_ (if applicable); and/or  
☐ is attached to this Declaration, Power of Attorney and Power to Inspect;

that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and

that I acknowledge my duty to disclose information which is material to the examination of this application in accordance with Rule 56(a) [37 C.F.R. §1.56(a)].

**CLAIM UNDER 35 U.S.C. §119:** I hereby claim foreign priority benefits under 35 U.S.C. §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application of which priority is claimed:

<u>Prior Foreign Application(s)</u> <u>Appln No.</u>	<u>Country</u>	<u>Filing Date</u> <u>Day-Mon-Year</u>	<u>Priority Claimed</u> <u>Yes - No</u>
9922071.7	Great Britain	17-09-1999	Yes

**POWER OF ATTORNEY:** As inventor, I hereby appoint **DANN, DORFMAN, HERRELL AND SKILLMAN, P.C.** of Philadelphia, Pennsylvania, and the following individual(s) as my attorneys or agents with full power of substitution to prosecute this application and to transact all business in the United States Patent and Trademark Office connected therewith: **Patrick J. Hagan, Reg. No. 27,643 and Kathleen D. Rigaut, Ph.D., Reg. 43,047.**

**POWER TO INSPECT:** I hereby give **DANN, DORFMAN, HERRELL AND SKILLMAN, P.C.** of Philadelphia, Pennsylvania or its duly accredited representatives power to inspect and obtain copies of the papers on file relating to this application.

**SEND CORRESPONDENCE TO:**

**CUSTOMER NUMBER 000110**

**DIRECT INQUIRIES TO:**

**Telephone: (215) 563-4100**  
**Facsimile: (215) 563-4044**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

**SOLE OR FIRST JOINT INVENTOR**


**SECOND JOINT INVENTOR (IF ANY)**

Full Name Caroline Dean  
First Middle Last  
Signature Caroline Dean  
Date 15 May 2002  
Residence Norwich UK GBX  
City State or Country  
Citizenship Brish  
Post Office Address:  
19 Waverley Rd  
Street Address  
Norwich Norfolk UK NR4 6SG  
City State or Country Zip Code

Full Name Yaron Yakov Levy  
First Middle Last  
Signature Yaron Levy  
Date April 19, 2002  
Residence ROSKILDE DENMARK DKX  
City State or Country  
Citizenship USA  
Post Office Address:  
VESTRE KIRKEVEJ 2G, 2TV  
Street Address  
ROSKILDE DENMARK DK-4000  
City State or Country Zip Code

10 ~~Month 33~~ 25 SEP 2002

#10



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of )  
 )  
Caroline Dean, et al. )  
 )  
Serial No. 10/088,187 ) Response to  
 ) Notification of Missing  
Filed: March 15, 2002 ) Requirements  
 )  
For: METHODS AND MEANS FOR )  
MODIFICATION OF PLANT FLOWERING )  
CHARACTERISTICS )

CERTIFICATE OF MAILING UNDER 37 C.F.R. §1.8(a)

I hereby certify that this correspondence is being deposited on September 19, 2002 with the United States Postal Service as first class mail in an envelope properly addressed to COMMISSIONER OF PATENTS AND TRADEMARKS, Washington, D.C. 20231.

September 19, 2002  
Date

Jane C. Bogan  
Jane C. Bogan

SUBMISSION OF SEQUENCE LISTING  
UNDER 37 C.F.R. §§1.821-1.825 AND PRELIMINARY AMENDMENT

The present submission is in response to a Notification of Missing Requirements dated July 19, 2002.

To comply with the requirements under 37 C.F.R. §§1.821-1.825, submitted herewith is a sequence listing of the amino acids presented in the above-referenced application. The sequence listing is being submitted in both paper copy and computer-readable form. Applicants respectfully request entry of the sequence listing into the above-identified patent application. The undersigned hereby verifies that the paper copy and computer readable form of the sequence listing are identical and do not contain any new matter.

In the event that a fee is required, the Commissioner is authorized to charge the account of the undersigned, Account No. 04-1406. A duplicate copy of this sheet is enclosed.

Respectfully submitted,  
DANN, DORFMAN, HERRELL AND SKILLMAN  
A Professional Corporation

By Patrick J. Hagan  
Patrick J. Hagan  
PTO Registration No. 27,643

Telephone: (215) 563-4100  
Facsimile: (215) 563-4044



10/088187 #6

10 ~~Aug 13 2002~~ 25 SEP 2002

1

SEQUENCE LISTING

<110> Dean, Caroline  
Levy, Yaron Y

<120> Methods and Means for Modification of Plant Flowering  
Characteristics

<130> 0380-P02825US0

<140> US 10/088,187

<141> 2002-03-15

<150> PCT/GB00/03525

<151> 2000-09-13

<150> GB 9922071.7

<151> 1999-09-17

<160> 48

<170> PatentIn Ver. 2.1

<210> 1

<211> 5000

<212> DNA

<213> Arabidopsis thaliana

<400> 1

tttaaaattc	gaattgggat	ttaagaaaaa	ttctcatcaa	atatttatca	ttagtgtata	60
tatatcagtg	ttttacattt	gttaatccta	aataataaac	cgatctgaaa	agttgataaa	120
tgcgttgtca	aaagacaaaa	tatacatcca	aacaaatcac	gtgattgcoct	tcaacttgcc	180
acgggttcaa	agattttaaca	aatcttctaa	aacaccaact	taaccacaga	atacacaagc	240
acagagtggg	ggtaaacata	caagttaatg	agttattcaa	atgagatttt	caatatcatt	300
cttcttcagc	ccgtcacaaag	aagccaagat	taagccatta	gaggaagttt	ataaacggac	360
aaaacctgct	tagatacaaa	gaatactagc	taatgtgttt	caacaaaact	caaattgacg	420
atacgttaca	ttcatattaa	tcacttcaga	gcttgattat	tcaaattatt	ttttctactg	480
tgatacatat	atacacacat	gttttgcttt	tctatgattc	tatctacatt	ttcataccgt	540
tgaataattt	atgtatgaat	tacgatgcaa	tttccctcat	tatgcttgaa	taaaatgctt	600
ttggacatgc	atgcgatatt	ggatctactt	ttggattota	tttttaaaaa	tcagcgagtt	660
tgttgctttg	taatttttaa	ttaggcataca	agaatttcta	aaatgcacgc	gaactggtga	720
aaagaggaat	gtttacgttt	acccctttat	tttcttacag	ctcataagga	tactgtcaga	780
agacagaacc	aaggctctct	gactataaat	tggaaatccat	ttaaacataa	tgttatgacc	840
aatgatggcc	aacgggttagc	ccaaaactaat	taactacaag	tcaagttcca	atattctaag	900
gagaaataat	agtatactaa	acatacatta	gagagggttaa	acttcttttt	ggatttaagt	960
gtgtatgcat	aggctattta	ttcttaagta	taactattaa	ctgtagctag	atttatacaa	1020
gaaatacata	aaactttatg	catgtgaggt	agccatgaat	atacgtacat	gttgcaatcg	1080
attatacatg	ttgtatttgg	atttctctat	acatgtttta	acttgctcatt	ctctaagtat	1140
atacatacca	ttaataactgt	gggcatgagt	ttatgataag	acttttcttt	tggagaccag	1200
ttttgttttc	ctttccacct	atatttgtct	ataggctcca	gacggtagac	tagttttacaa	1260
gtgtttttat	atgtttctaaa	taaaattgag	attttccgga	acggtatgat	ctgtttgcaa	1320
ataaggacgt	atatataaca	gtatcaaaata	tatttggtgt	tataaggcaa	taatatattt	1380
tctgagatat	tgcgtgttac	aaaaaagaaa	tatttggtta	gaaaaaaaaa	gatgggtcgaa	1440
aaaggggagt	aggtgggggc	ggtcggcttt	tgattagtta	ataaaagaaa	ccacacgagt	1500
gacctaccga	ttcgactcaa	cgagtctacc	gagctaacac	agattoaaact	cgctcgagct	1560
tcgtttttatg	acaagttggt	tttttttttt	tttttttaat	tttttcatct	tcttgggttt	1620
ggttggttca	ctcttcaggt	caggtgtgta	aaaaagaaag	aaagaaaaga	gagattgttg	1680
tgttgtaacc	cctttgacta	aaatctaata	aactttttta	acacaacaaa	actccttcag	1740
atctgaaagg	gttcttcttc	tctcttagtc	tctttgtcct	tttattctcc	gtcgtcgttt	1800
catgatctga	ctctctggtc	ttctcttctt	cttctctctc	ttctattttt	tcttacttcg	1860

toactgttgt	gtctgaacat	gccacgccct	ttcttccata	agttgatttt	ctcatccaact	1920
atccaagaaa	aacgtctggg	aacttactct	ctctctctct	ctctctctct	gttctccttc	1980
toctcatctt	tcaaagtttt	gattttgtgc	gaaattgagg	gttttcaagg	tttggaaatct	2040
ggtgaacgag	tttgaagat	tatgccttgt	gacactcttg	cttgatttct	tacaattcac	2100
ttgtattgat	tctttgtaag	aatcgagtc	aggttgtgct	tttatcttct	tactcttccc	2160
tgttttgggt	aatgaaaaga	agttccattt	ttgaactttg	tgttgtotta	ttggtaaaat	2220
gagaatttgt	gggtttccaa	tggaagtctg	caagacagtt	tcttttggtc	attgggtgtg	2280
tttgggtggg	aattgggtat	ttgatgggat	atctgtactc	tgacagcata	ttgtgtgtag	2340
tttgggaatt	tttttttttt	ttttgagtga	tttgactttt	ggaggacgat	ttgattctgt	2400
cagattgata	aaatttcttc	tgaggagaaa	aagttgagat	ctgtttatgg	tttctctatt	2460
ataaatgtct	gttttgttta	ctctattttg	actgttttct	ctgtttgact	taggaatgtc	2520
tgagatctta	gactccttat	tgagtattgt	gtggcttgtg	agtgaatccc	taaaactgag	2580
tagttgactt	gttttgaagg	tctctatgta	ttgtgtctat	gttttaaagt	tgtctacttt	2640
atgtgataca	gtgattagtc	atcacttgta	cagattcccc	caagagcatt	gttttgaaca	2700
aatccaaatt	tgcttagctc	tccatttggt	atttaagtga	ctagattttc	tctggaataa	2760
tgatttcgat	taacacaggc	atattatgtg	aaccaagttt	gcaaattatt	aatgtgataa	2820
gatcatagga	gtcgtgtaat	caatctattc	agagataaat	gtaccatttt	acatgtgtac	2880
taatggactg	tgtctccttg	ttgatgcctt	ctctaaactg	aaatatggcc	ttttggtttg	2940
tgtttttaaa	ttaggtaaaag	ccgtcgtttc	ttcagctact	gtgtttattg	gatgtttttg	3000
ctgaaaaatg	tctgtttcga	tttgatgttc	tcgcaatatt	ctgtgctgtt	cttatagata	3060
ttgtggacat	ttatatcatt	atatgcttct	ttatatctca	taccggcatg	cttgtgcaga	3120
gggtcccaga	taagtttgtg	agtaaatcca	aggatgagct	ttcggttgct	gttgcaactc	3180
cagtacctga	tggtcatggt	tgrcgtgtag	gactaaggaa	agctgacaac	aaaatttggg	3240
ttcaagatgg	ttggcaagag	tttgttgacc	gttactccat	tcgactgggt	tatcttttga	3300
tttttagata	tgaaggaaac	tctgccttca	cgctctacat	tttcaattta	tcccactctg	3360
agatcaatta	ccattccacc	ggtctcatgg	attccgctca	caaccaactc	aaacgcgccc	3420
gtttgtttga	agaccttgaa	gatgaagatg	ccgagggtcat	ctttccttct	tctgtgtacc	3480
catcaccact	tcctgagctc	acagtaccag	ccaacaaagg	gtatgctagt	tcagccatcc	3540
aaacctgtgt	cactggacca	gttaaaagtg	atatttataa	ccaactgatt	ccctttatct	3600
atcgctgatt	acgcgtctta	tcattctttt	gaggttgatg	cttgatattt	tccttatctc	3660
cagctgaaga	gccaaaccca	accccaaaaa	tacctaaaaa	gagagggagg	aagaagaaaa	3720
atgctgatcc	tggttaagcac	ttttcctctt	tgaaatgctt	cagactcggt	ttcagaggat	3780
tcacagattc	ttcctcatga	tacatatatc	cttttgatat	tgctccttaca	gaggaaataa	3840
actcatcagc	tccgcgagat	gatgatccag	agaaccgttc	aaagttctac	gagagtgtct	3900
ctgcgagaaa	gagaaccgtg	actgcagaag	aaagagagag	agccatcaat	gcagccaaaa	3960
cgttcgaacc	aacaaaccc	ttcttcagag	tggttctgct	accatcctat	ctatacagag	4020
gttgcatcat	ggtaataaaa	aaacatotta	ggaagactta	atcttatcgg	tgtcttcaat	4080
gatctctaaa	agaagccttc	tgtttctgtt	tctctcaaca	gtatcttctc	tctgggtttg	4140
ctgagaagta	cctaagtggg	atctccgggt	tcatacaaagt	ccagcttgct	gagaaacaat	4200
ggcctgttcg	atgtctctac	aaagccggga	gagccaaatt	cagtcaagga	tggtacgaat	4260
tcactctaga	gaacaactta	ggagaaggag	acgtctgtgt	gtttgagctg	ctcagaacca	4320
gagatttctg	tttgaaagtg	acagcctttc	gagtcacaag	gtacgtctga	acaaagcatt	4380
atgggtgtgat	cattctggat	ttgcaagtac	aatgtogtgt	aggagtatct	taatttataa	4440
acaactaaaa	aactctcttc	tggtctgtgt	cattattgctg	tcagtgtctc	gttttttctc	4500
tcgggtttac	tttgtgttat	cgatgtggat	aagttgtttt	tacctatta	tatataacct	4560
cttgagtggg	actcaaattg	tttgagttag	acaaacaaag	ttagggttta	agaagaagtc	4620
tgtaaatacc	taatctccat	caaattttgag	tagaaagaca	aactgttctg	gtggaatata	4680
aggaggggac	ttgagataac	aaacttaaga	atagccttca	agccaacgtc	tagaattttg	4740
tgaagttgtt	gtttgatcac	ctctgagata	attggaacac	ctcttcatgc	agtttgcttg	4800
aggatactgg	tgaaaaatggg	agtattgaag	gaaaatgcac	atataagatt	gtaggtggga	4860
actgtggtag	cagacacaac	actgtttctc	tagacatata	ctgtaccaga	catgtttttg	4920
tcataaaaact	taaaaaaaag	aaaaccgtgt	gtaaatcaag	caaggaacaa	ctacaatatt	4980
acaatcttat	tgagatatca					5000

```
<400> 5
Asp Gly His Val
1
```

<400> 9  
Lys Lys Met Leu Ile Leu Arg Lys  
1 5

```
<210> 10
<211> 1495
<212> DNA
<213> Arabidopsis thaliana
```

<400> 10																			
tcttggggttt	ggttgggtcoa	ctcttcaggt	caggtgtgta	aaaaagaaag	aaagaaaaga	60													
gagattgttg	tgttgtaacc	cctttgacta	aaatctaatt	aactttttta	acacaacaaa	120													
actccttcag	atctgaaagg	gttctttctt	tctcttagtc	tctttgtcct	tttattctcc	180													
gtcgtcgttt	catgatctga	ctctctggtc	ttctcttctt	cttctttctt	ttctattttt	240													
tcttacttcg						tcactgttgt	gtctgaac	atg	cca	cgc	cct	ttc	ttc	cat	aag	292			
								Met	Pro	Arg	Pro	Phe	Phe	His	Lys				
								1							5				
ttg	att	ttc	tca	tcc	act	atc	caa	gaa	aaa	cgt	ctg	agg	gtc	cca	gat	340			
Leu	Ile	Phe	Ser	Ser	Thr	Ile	Gln	Glu	Lys	Arg	Leu	Arg	Val	Pro	Asp				
						10							15						
aag	ttt	gtg	agt	aaa	ttc	aag	gat	gag	ctt	tgc	gtt	gct	gtt	gca	ctc	388			
Lys	Phe	Val	Ser	Lys	Phe	Lys	Asp	Glu	Leu	Ser	Val	Ala	Val	Ala	Leu				
						25							30						
aca	gta	cct	gat	ggg	cat	gtt	tgg	cgt	gta	gga	cta	agg	aaa	gct	gac	436			
Thr	Val	Pro	Asp	Gly	His	Val	Trp	Arg	Val	Gly	Leu	Arg	Lys	Ala	Asp				
						45							50						
aac	aaa	att	tgg	ttt	caa	gat	ggg	tgg	caa	gag	ttt	gtt	gac	cgt	tac	484			
Asn	Lys	Ile	Trp	Phe	Gln	Asp	Gly	Trp	Gln	Glu	Phe	Val	Asp	Arg	Tyr				
						60							65						
tcc	att	cgc	att	ggg	tat	ctt	ttg	att	ttt	aga	tat	gaa	gga	aac	tct	532			
Ser	Ile	Arg	Ile	Gly	Tyr	Leu	Leu	Ile	Phe	Arg	Tyr	Glu	Gly	Asn	Ser				
						75							80						
gcc	ttc	agc	gtc	tac	att	ttc	aat	tta	tcc	cac	tct	gag	atc	aat	tac	580			
Ala	Phe	Ser	Val	Tyr	Ile	Phe	Asn	Leu	Ser	His	Ser	Glu	Ile	Asn	Tyr				
						90							95						
cat	tcc	acc	ggg	ctc	atg	gat	tcc	gct	cac	aac	cac	ttc	aaa	cgc	gcc	628			
His	Ser	Thr	Gly	Leu	Met	Asp	Ser	Ala	His	Asn	His	Phe	Lys	Arg	Ala				
						105							110						
cgt	ttg	ttt	gaa	gac	ctt	gaa	gat	gaa	gat	gcc	gag	gtc	atc	ttt	cct	676			
Arg	Leu	Phe	Glu	Asp	Leu	Glu	Asp	Glu	Asp	Ala	Glu	Val	Ile	Phe	Pro				
						125							130						
tct	tct	gtg	tac	cca	tca	cca	ctt	cct	gag	tct	aca	gta	cca	gcc	aac	724			
Ser	Ser	Val	Tyr	Pro	Ser	Pro	Leu	Pro	Glu	Ser	Thr	Val	Pro	Ala	Asn				
						140							145						
aaa	ggg	tat	gct	agt	tca	gcc	atc	caa	acc	ttg	ttc	act	gga	cca	gtt	772			
Lys	Gly	Tyr	Ala	Ser	Ser	Ala	Ile	Gln	Thr	Leu	Phe	Thr	Gly	Pro	Val				
						155							160						

<400> 11  
Met Pro Arg Pro Phe Phe His Lys Leu Ile Phe Ser Ser Thr Ile Gln  
1 5 10 15



Glu Lys Arg Leu Arg Val Pro Asp Lys Phe Val Ser Lys Phe Lys Asp  
 20 25 30  
 Glu Leu Ser Val Ala Val Ala Leu Thr Val Pro Asp Gly His Val Trp  
 35 40 45  
 Arg Val Gly Leu Arg Lys Ala Asp Asn Lys Ile Trp Phe Gln Asp Gly  
 50 55 60  
 Trp Gln Glu Phe Val Asp Arg Tyr Ser Ile Arg Ile Gly Tyr Leu Leu  
 65 70 75 80  
 Ile Phe Arg Tyr Glu Gly Asn Ser Ala Phe Ser Val Tyr Ile Phe Asn  
 85 90 95  
 Leu Ser His Ser Glu Ile Asn Tyr His Ser Thr Gly Leu Met Asp Ser  
 100 105 110  
 Ala His Asn His Phe Lys Arg Ala Arg Leu Phe Glu Asp Leu Glu Asp  
 115 120 125  
 Glu Asp Ala Glu Val Ile Phe Pro Ser Ser Val Tyr Pro Ser Pro Leu  
 130 135 140  
 Pro Glu Ser Thr Val Pro Ala Asn Lys Gly Tyr Ala Ser Ser Ala Ile  
 145 150 155 160  
 Gln Thr Leu Phe Thr Gly Pro Val Lys Ala Glu Glu Pro Thr Pro Thr  
 165 170 175  
 Pro Lys Ile Pro Lys Lys Arg Gly Arg Lys Lys Lys Asn Ala Asp Pro  
 180 185 190  
 Glu Glu Ile Asn Ser Ser Ala Pro Arg Asp Asp Asp Pro Glu Asn Arg  
 195 200 205  
 Ser Lys Phe Tyr Glu Ser Ala Ser Ala Arg Lys Arg Thr Val Thr Ala  
 210 215 220  
 Glu Glu Arg Glu Arg Ala Ile Asn Ala Ala Lys Thr Phe Glu Pro Thr  
 225 230 235 240  
 Asn Pro Phe Phe Arg Val Val Leu Arg Pro Ser Tyr Leu Tyr Arg Gly  
 245 250 255  
 Cys Ile Met Tyr Leu Pro Ser Gly Phe Ala Glu Lys Tyr Leu Ser Gly  
 260 265 270  
 Ile Ser Gly Phe Ile Lys Val Gln Leu Ala Glu Lys Gln Trp Pro Val  
 275 280 285  
 Arg Cys Leu Tyr Lys Ala Gly Arg Ala Lys Phe Ser Gln Gly Trp Tyr  
 290 295 300  
 Glu Phe Thr Leu Glu Asn Asn Leu Gly Glu Gly Asp Val Cys Val Phe  
 305 310 315 320  
 Glu Leu Leu Arg Thr Arg Asp Phe Val Leu Lys Val Thr Ala Phe Arg  
 325 330 335  
 Val Asn Glu Tyr Val  
 340

<210> 12

<211> 1495

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: vrn1-1  
 mutation

<400> 12

tcttgggttt ggttgggtca ctcttcaggt caggtgtgta aaaaagaaag aaagaaaaga 60  
 gagattgttg tgttgtaacc cctttgacta aaatctaata aactttttta acacaacaaa 120  
 actccttcag atctgaaagg gttcttcttc tctcttagtc tctttgtcct tttattctcc 180  
 gtcgtcgttt catgatctga ctctctgggc ttctcttctt cttctctctc ttctattttt 240  
 tcttacttcg tcaactgttg gtctgaacat gccacgccct ttcttcata agttgatttt 300  
 ctcatccact atccaagaaa aacgtctgag ggtcccagat aagtttgtga gtaaattcaa 360

8

```

ggatgagctt tgggttgcgt ttgcactcac agtacctgat ggtcatgttt gacgtgtagg 420
actaaggaaa gctgacaaca aaattttggt tcaagatggt tggcaagagt ttgttgaccg 480
ttactccatt cgcattgggt atcttttgat ttttagatat gaaggaaact ctgccttcag 540
cgtctacatt ttcaatttat ccactctga gatcaattac cattccaccg gtctcatgga 600
ttccgctcac aaccacttca aacgcgcccg tttgtttgaa gaccttgaag atgaagatgc 660
cgaggtcatc tttccttctt ctgtgtaccc atcaccactt cctgagtcta cagtaccagc 720
caacaaaggg tatgctagtt cagccatcca aaccttggtc actggaccag ttaaagctga 780
agagccaacg ccaaccccaa aaatacctaa aaagagaggg aggaagaaga aaaatgctga 840
tcctgaggaa ataaactcat cagctccgcg agatgatgat ccagagaacc gttcaaagtt 900
ctacgagagt gcttctgcga gaaagagaac cgtgactgca gaagaaagag agagagccat 960
caatgcagcc aaaacgttcg aaccaacaaa ccctttcttc agagtgggtc tgcgaccatc 1020
ctatctatac agaggttgca tcatgtatct tccttctggg tttgctgaga agtacctaag 1080
tggtatctcc gggttcatca aagtccagct tgcggagaaa caatggcctg ttcatgtctt 1140
ctacaaagcc gggagagcca aattcagtca aggatggtac gaattcactc tagagaacaa 1200
cttaggagaa ggagacgtct gtgtgtttga gctgctcaga accagagatt tcgttttgaa 1260
agtgcagacc tttcgagtca acgagtacgt ctgaacaaaag cattatggtg tgatcattct 1320
ggatttgcaa gtacaatgtc gtgtaggagt atcttaattt aaaaaaactt aaaaaactct 1380
cttctgggtc gtgtcattat tgcgtcagtg tctcgttttt tctctcgggt ttactttgtg 1440
ttatcgatgt ggataagttg tttttacctc attatatata acctcttgag tggaa 1495

```

<210> 13

<211> 1494

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: vrn1-2  
mutation

<400> 13

```

tcttgggttt ggttgggtca ctcttcaggt caggtgtgta aaaaagaaag aaagaaaaga 60
gagattgttg tgttgaacc ctttgacta aaatctaata aactttttta acacaacaaa 120
actccttcag atctgaaagg gttctcttct tctcttagtc tctttgtcct tttattctoc 180
gtcgtcgttt catgatctga ctctctggtc ttctcttctt ctctctcttc ttctattttt 240
tcttacttcg tcaactgttg gtctgaacat gccacgccct ttcttccata agttgatttt 300
ctcatccact atccaagaaa aacgtctgag ggtcccagat aagtttgtga gtaaattcaa 360
ggatgagctt tgggttgcgt ttgcactcac agtacctgat ggtcatgttt ggcgtgtagg 420
actaaggaaa gctgacaaca aaattttggt tcaagatggt tggcaagagt ttgttgaccg 480
ttactccatt cgcattgggt atcttttgat ttttagatat gaaggaaact ctgccttcag 540
cgtctacatt ttcaatttat ccactctga gatcaattac cattccaccg gtctcatgga 600
ttccgctcac aaccacttca aacgcgcccg tttgtttgaa gaccttgaag atgaagatgc 660
cgaggtcatc tttccttctt ctgtgtaccc atcaccactt cctgagtcta cagtaccagc 720
caacaaaggg tatgctagtt cagccatcca aaccttggtc actggaccag ttaaagctga 780
agagccaacg ccaaccccaa aaatacctaa aaagagaggg aggaagaaga aaatgctgat 840
cctgaggaaa taaactcatc agctccgcga gatgatgatc cagagaaccg ttcaaagttc 900
tacgagagtg cttctgcgag aaagagaacc gtgactgcag aagaaagaga gagagccatc 960
aatgcagcca aaacgttcga accaacaac cctttcttca gagtggttct gcgaccatcc 1020
tatctataca gaggttgcat catgtatctt ccttctgggt ttgctgagaa gtacctaagt 1080
gggatctccg ggttcacaa agtccagctt cgggagaaac aatggcctgt tcgatgtctc 1140
tacaagcccg ggagagccaa attcagtcaa ggtggttacg aattcactct agagaacaac 1200
ttaggagaag gagacgtctg tgtgtttgag ctgctcagaa ccagagattt cgttttgaaa 1260
gtgacagcct ttcgagtcaa cgagtacgtc tgaacaaagc attatggtgt gatcattctg 1320
gatttgcaag tacaatgtcg tgtaggagta tcttaattta aaaacaacta aaaaaactct 1380
ttctgggtct gtgcattatt gcgtcagtg ctctgttttt ctctcgggtt tactttgtgt 1440
tatogatgtg gataagttgt ttttacctca ttatatataa cctcttgagt ggaa 1494

```

<210> 14  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 14  
 acctgcttct gccaacgct c

21

<210> 15  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 15  
 agttcgctct tgctgttttt ttcccc

26

<210> 16  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 16  
 cctcttcgct attacgccag

20

<210> 17  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 17  
 gcccttccca acagttcg

18

<210> 18  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 18  
 cacacaggaa acagctat

18

<210> 19  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 19  
 acacaacata cgagccggaa 20

<210> 20  
 <211> 17  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 20  
 caacggttag cccaaac 17

<210> 21  
 <211> 17  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 21  
 gtttgggcta accggtg 17

<210> 22  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 22  
 gagaccagtt ttgttttcc 19

<210> 23  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 23  
gacaaatata ggtggaaagg 20

<210> 24  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 24  
aaaggggagt aggtggg 17

<210> 25  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 25  
ctctctggtc ttctcttc 18

<210> 26  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 26  
gaagagaaga ccagagag 18

<210> 27  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 27  
ttttctcatc cactatcc 18

<210> 28  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 28  
 tttcttggat agtggatgag 20

<210> 29  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 29  
 aaaacaggga agagtaagaa g 21

<210> 30  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 30  
 cattggttgt gtttggtggg 20

<210> 31  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 31  
 ggtctctatg tattgtgc 18

<210> 32  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

18

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide



<400> 41  
ctctctctct ttctttctg 18

<210> 42  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 42  
ccactctgaa gaaaggg 17

<210> 43  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 43  
ccttctgttt ctgtttctc 19

<210> 44  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 44  
gagaaacaga aacagaagg 19

<210> 45  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 45  
aagatactcc tacacgac 18

<210> 46  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 46  
 gtctcgtttt ttctctcgg

19

<210> 47  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 47  
 ctaccacagt tcccacctac


20

<210> 48  
 <211> 301  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 48  
 Met Pro Arg Ser Phe Phe His Met Phe Asn Ser Leu Phe Leu Ser Ser  
 1 5 10 15  
 Thr Gln Ala Ser Gly Leu Arg Lys Ala Asn Asn Lys Ile Trp Phe Gln  
 20 25 30  
 Asp Gly Trp Gln Glu Phe Val Asn Arg Phe Ser Ile Arg Ile Gly Phe  
 35 40 45  
 Arg Tyr Lys Val Thr Val Tyr Ile Phe Gln Phe Tyr Pro Pro His Ser  
 50 55 60  
 Glu Ile Asn His His Ser Ser Ser Glu Ala Leu Met Gln Met Asp Ser  
 65 70 75 80  
 Ala Gln Asn Gln Phe Asn Lys Arg Ala Arg Leu Phe Glu Asp Pro Glu  
 85 90 95  
 Leu Lys Asp Ala Lys Val Ile Tyr Pro Ser Asn Pro Glu Ser Thr Glu  
 100 105 110  
 Pro Val Asn Lys Gly Tyr Gly Gly Ser Thr Ala Ile Gln Ser Phe Phe  
 115 120 125  
 Lys Glu Ser Lys Ala Glu Glu Thr Pro Lys Val Leu Lys Lys Arg Gly  
 130 135 140

Arg	Lys	Lys	Lys	Asn	Pro	Asn	Pro	Glu	Glu	Val	Asn	Ser	Ser	Thr	Pro	145	150	155	160
Gly	Gly	Asp	Asp	Ser	Glu	Asn	Arg	Ser	Lys	Phe	Tyr	Glu	Ser	Ala	Ser	165	170	175	
Ala	Arg	Lys	Arg	Thr	Val	Thr	Ala	Glu	Glu	Arg	Glu	Arg	Ala	Val	Asn	180	185	190	
Ala	Ala	Lys	Thr	Phe	Glu	Pro	Thr	Asn	Pro	Tyr	Phe	Arg	Val	Val	Leu	195	200	205	
Arg	Pro	Ser	Tyr	Leu	Tyr	Arg	Gly	Cys	Ile	Met	Tyr	Leu	Pro	Ser	Gly	210	215	220	
Phe	Ala	Glu	Lys	Tyr	Leu	Ser	Gly	Ile	Ser	Gly	Phe	Ile	Lys	Leu	Gln	225	230	235	240
Leu	Gly	Glu	Lys	Gln	Trp	Pro	Val	Arg	Cys	Leu	Tyr	Lys	Ala	Gly	Arg	245	250	255	
Ala	Lys	Phe	Ser	Gln	Gly	Trp	Tyr	Glu	Phe	Thr	Leu	Glu	Asn	Asn	Ile	260	265	270	
Gly	Glu	Gly	Asp	Val	Cys	Val	Phe	Glu	Leu	Leu	Arg	Thr	Arg	Asp	Phe	275	280	285	
Val	Leu	Glu	Val	Thr	Ala	Phe	Arg	Val	Asn	Glu	Tyr	Val				290	295	300	

United States Patent & Trademark Office  
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not  
present  
for scanning. (Document title)

✓ Scanned copy is best available. Some drawings are dark.